

Physics 741 – Graduate Quantum Mechanics 1
Solutions to Chapter 3

3.5 [5] Prove the parity operator Π , defined by (3.40) is both Hermitian and unitary.

To show it is Hermitian, we must show that $\langle \phi | \Pi | \psi \rangle^* = \langle \psi | \Pi | \phi \rangle$, so

$$\langle \phi | \Pi | \psi \rangle^* = \left[\int d^3\mathbf{r} \phi^*(\mathbf{r}) \psi(-\mathbf{r}) \right]^* = \int d^3\mathbf{r} \psi^*(-\mathbf{r}) \phi(\mathbf{r}) = \int d^3\mathbf{r} \psi^*(\mathbf{r}) \phi(-\mathbf{r}) = \langle \psi | \Pi | \phi \rangle$$

Now that we know it is Hermitian, we can take advantage of this to show that

$$\Pi^\dagger \Pi \psi(\mathbf{r}) = \Pi^2 \psi(\mathbf{r}) = \Pi \psi(-\mathbf{r}) = \psi(\mathbf{r}).$$

Since this is true for all wave functions, it follows that $\Pi^\dagger \Pi = 1$.

3.6 [15] Consider the Hermitian matrix: $H = E_0$

$$H = E_0 \begin{pmatrix} 0 & 3i & 0 \\ -3i & 8 & 0 \\ 0 & 0 & 8 \end{pmatrix}$$

(a) [10] Find all three eigenvalues and eigenvectors of H .

Removing the common factor of E_0 we note that H is block-diagonal, as I have sketched in with dashed lines in the problem itself, reducing the matrix to a 2×2 matrix and a trivial 1×1 matrix:

$$H_2 = \begin{pmatrix} 0 & 3i \\ -3i & 8 \end{pmatrix} \quad \text{and} \quad H_1 = (8)$$

The matrix H_1 has eigenvalue 8, and eigenvector (1), which makes it trivial. The eigenvalues of matrix H_2 can be found using the characteristic equation

$$0 = \det(H_2 - \lambda \mathbf{1}) = \begin{vmatrix} -\lambda & 3i \\ -3i & 8 - \lambda \end{vmatrix} = \lambda^2 - 8\lambda - (3i)(-3i) = \lambda^2 - 8\lambda - 9 = (\lambda - 9)(\lambda + 1)$$

This has solutions $\lambda = 9$ and $\lambda = -1$. To find each of these values, we put in an arbitrary vector and solve the eigenvalue equation. For example, for $\lambda = 9$, we have

$$\begin{pmatrix} 0 & 3i \\ -3i & 8 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = 9 \begin{pmatrix} \alpha \\ \beta \end{pmatrix},$$

$$\begin{pmatrix} 3i\beta \\ -3i\alpha + 8\beta \end{pmatrix} = \begin{pmatrix} 9\alpha \\ 9\beta \end{pmatrix}.$$

The first of these equations implies $\beta = -3i\alpha$; if we plug this into the second, we find that it is also automatically satisfied. We also want the eigenvector normalized, so

$$1 = |\alpha|^2 + |\beta|^2 = 10|\alpha|^2$$

We have an arbitrary phase to choose; if we pick α to be real and positive, $\alpha = 1/\sqrt{10}$, and we have the eigenvector

$$|9\rangle = \frac{1}{\sqrt{10}} \begin{pmatrix} 1 \\ -3i \end{pmatrix}$$

For the other eigenvector, we have

$$\begin{pmatrix} 0 & 3i \\ -3i & 8 \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix} = - \begin{pmatrix} \alpha \\ \beta \end{pmatrix},$$

$$\begin{pmatrix} 3i\beta \\ -3i\alpha + 8\beta \end{pmatrix} = \begin{pmatrix} -\alpha \\ -\beta \end{pmatrix}$$

Both equations imply $\beta = i\alpha/3$. Our normalization condition becomes

$$1 = |\alpha|^2 + |\beta|^2 = \frac{10}{9}|\alpha|^2$$

Once again we pick α to be real and positive, $\alpha = 3/\sqrt{10}$, and we have the eigenvector

$$|-1\rangle = \frac{1}{\sqrt{10}} \begin{pmatrix} 3 \\ i \end{pmatrix}$$

Returning to the full three-dimensional space and restoring E_0 , our eigenvectors are

$$|-E_0\rangle = \frac{1}{\sqrt{10}} \begin{pmatrix} 3 \\ i \\ 0 \end{pmatrix}, \quad |9E_0\rangle = \frac{1}{\sqrt{10}} \begin{pmatrix} 1 \\ -3i \\ 0 \end{pmatrix}, \quad |8E_0\rangle = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

Your answers might be slightly different, in that the phases could be different, or the eigenvectors could be listed in a different order.

(b) [5] Construct the unitary matrix V which diagonalizes H . Check explicitly that $V^\dagger V = 1$ and $V^\dagger H V = H'$ is real and diagonal.

The unitary matrix V just consists of the eigenvectors listed in any order, so we have

$$V = \begin{pmatrix} \frac{3}{\sqrt{10}} & \frac{1}{\sqrt{10}} & 0 \\ \frac{i}{\sqrt{10}} & -\frac{3i}{\sqrt{10}} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Your answer could be different, in that the columns could come in a different order, and each column could be multiplied by an arbitrary phase.

We have ahead of us some boring matrix multiplication.

$$\begin{aligned}
V^\dagger V &= \begin{pmatrix} \frac{3}{\sqrt{10}} & -\frac{i}{\sqrt{10}} & 0 \\ \frac{1}{\sqrt{10}} & \frac{3i}{\sqrt{10}} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \frac{3}{\sqrt{10}} & \frac{1}{\sqrt{10}} & 0 \\ \frac{i}{\sqrt{10}} & -\frac{3i}{\sqrt{10}} & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} \frac{9}{10} + \frac{1}{10} & \frac{3}{10} - \frac{3}{10} & 0 \\ \frac{3}{10} - \frac{3}{10} & \frac{1}{10} + \frac{9}{10} & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
V^\dagger H V &= E_0 \begin{pmatrix} \frac{3}{\sqrt{10}} & -\frac{i}{\sqrt{10}} & 0 \\ \frac{1}{\sqrt{10}} & \frac{3i}{\sqrt{10}} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 3i & 0 \\ -3i & 8 & 0 \\ 0 & 0 & 8 \end{pmatrix} \begin{pmatrix} \frac{3}{\sqrt{10}} & \frac{1}{\sqrt{10}} & 0 \\ \frac{i}{\sqrt{10}} & -\frac{3i}{\sqrt{10}} & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
&= E_0 \begin{pmatrix} \frac{3}{\sqrt{10}} & -\frac{i}{\sqrt{10}} & 0 \\ \frac{1}{\sqrt{10}} & \frac{3i}{\sqrt{10}} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} -\frac{3}{\sqrt{10}} & \frac{9}{\sqrt{10}} & 0 \\ -\frac{i}{\sqrt{10}} & -\frac{27i}{\sqrt{10}} & 0 \\ 0 & 0 & 8 \end{pmatrix} = E_0 \begin{pmatrix} -\frac{9}{10} - \frac{1}{10} & \frac{27}{10} - \frac{27}{10} & 0 \\ -\frac{3}{10} + \frac{3}{10} & \frac{9}{10} + \frac{81}{10} & 0 \\ 0 & 0 & 8 \end{pmatrix} = E_0 \begin{pmatrix} -1 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 8 \end{pmatrix}
\end{aligned}$$

As you can see, $V^\dagger V = 1$ and $V^\dagger H V$ is real and diagonal (and has the eigenvalues on its diagonal).