

PHY 711 – Notes on Hydrodynamics – (“Solitary Waves”[1])

Basic assumptions

We assume that we have in incompressible fluid ($\rho = \text{constant}$) a velocity potential of the form $\Phi(x, z, t)$. The surface of the fluid is described by $h + \zeta(x, t) = z$. The fluid is contained in a tank with a structureless bottom (defined by the plane $z = 0$) and is filled to a vertical height h at equilibrium. These functions satisfy the following conditions.

Poisson equation:

$$\frac{\partial^2 \Phi(x, z, t)}{\partial x^2} + \frac{\partial^2 \Phi(x, z, t)}{\partial z^2} = 0 \quad (1)$$

Zero vertical velocity at bottom of the tank:

$$\frac{\partial \Phi(x, 0, t)}{\partial z} = 0 \quad (2)$$

Bernoulli’s equation:

$$-\frac{\partial \Phi(x, z, t)}{\partial t} + \frac{1}{2} \left[\left(\frac{\partial \Phi(x, z, t)}{\partial x} \right)^2 + \left(\frac{\partial \Phi(x, z, t)}{\partial z} \right)^2 \right] + g\zeta(x, t) \Big|_{z=h+\zeta} = 0 \quad (3)$$

Surface equation:

$$-\frac{\partial \Phi(x, z, t)}{\partial z} + \frac{\partial \Phi(x, z, t)}{\partial x} \frac{\partial \zeta(x, t)}{\partial x} - \frac{\partial \zeta(x, t)}{\partial t} \Big|_{z=h+\zeta} = 0 \quad (4)$$

In this treatment, we assume seek the form of surface waves traveling along the x - direction and assume that the effective wavelength is much larger than the height of the surface h . This allows us to approximate the z - dependence of $\Phi(x, z, t)$ by means of a Taylor series expansion:

$$\Phi(x, z, t) \approx \Phi(x, 0, t) + z \frac{\partial \Phi}{\partial z}(x, 0, t) + \frac{z^2}{2} \frac{\partial^2 \Phi}{\partial z^2}(x, 0, t) + \frac{z^3}{3!} \frac{\partial^3 \Phi}{\partial z^3}(x, 0, t) + \frac{z^4}{4!} \frac{\partial^4 \Phi}{\partial z^4}(x, 0, t) \cdots \quad (5)$$

This expansion can be simplified because of the bottom boundary condition (2) which ensures that all odd derivatives $\frac{\partial^n \Phi}{\partial z^n}(x, 0, t)$ vanish from the Taylor expansion (5). In addition, the Poisson equation (1) allows us to convert all even derivatives with respect to z to derivatives with respect to x . Therefore, the expansion (5) becomes:

$$\Phi(x, z, t) \approx \Phi(x, 0, t) - \frac{z^2}{2} \frac{\partial^2 \Phi}{\partial x^2}(x, 0, t) + \frac{z^4}{4!} \frac{\partial^4 \Phi}{\partial x^4}(x, 0, t) \cdots \quad (6)$$

For convenience we define $\phi(x, t) \equiv \Phi(x, 0, t)$. Using Eq. (6), the Bernoulli equation (3) then becomes:

$$-\frac{\partial \phi}{\partial t} + \frac{(h + \zeta)^2}{2} \frac{\partial^3 \phi}{\partial t \partial x^2} + \frac{1}{2} \left[\left(\frac{\partial \phi}{\partial x} \right)^2 + \left((h + \zeta) \frac{\partial^2 \phi}{\partial x^2} \right)^2 \right] + g\zeta = 0, \quad (7)$$

where we have discarded some of the higher order terms. Keeping all terms up to leading order in non-linearity and up to fourth order derivatives in the linear terms, the Bernoulli equation becomes:

$$-\frac{\partial\phi}{\partial t} + \frac{h^2}{2} \frac{\partial^3\phi}{\partial t\partial x^2} + \frac{1}{2} \left(\frac{\partial\phi}{\partial x}\right)^2 + g\zeta = 0. \quad (8)$$

Using a similar analysis and approximation, the surface definition equation (4) becomes:

$$\frac{\partial}{\partial x} \left((h + \zeta(x, t)) \frac{\partial\phi}{\partial x} \right) - \frac{h^3}{3!} \frac{\partial^4\phi}{\partial x^4} - \frac{\partial\zeta}{\partial t} = 0, \quad (9)$$

We would like to solve Eqs. (8-9) for a traveling wave of the form:

$$\phi(x, t) = \chi(x - ct) \text{ and } \zeta(x, t) = \eta(x - ct), \quad (10)$$

where the speed of the wave c will be determined. Letting $u \equiv x - ct$, Eqs. (8 and 9) become:

$$\frac{d}{du} \left((h + \eta(u)) \frac{d\chi(u)}{du} \right) - \frac{h^3}{6} \frac{d^4\chi(u)}{du^4} + c \frac{d\eta(u)}{du} = 0, \quad (11)$$

and

$$c \frac{d\chi(u)}{du} - \frac{ch^2}{2} \frac{d^3\chi(u)}{du^3} + \frac{1}{2} \left(\frac{d\chi(u)}{du} \right)^2 + g\eta(u) = 0. \quad (12)$$

The modified surface equation (11) can be integrated once with respect to u , choosing the constant of integration to be zero and giving the new form for the surface condition:

$$(h + \eta)\chi' - \frac{h^3}{6}\chi''' + c\eta = 0, \quad (13)$$

where we have abbreviated derivatives with respect to u with the “'” symbol. This equation, and the modified Bernoulli equation (8) are now two coupled non-linear equations. In order to solve them, we use, the modified Bernoulli equation to approximate $\chi'(u)$ and its higher derivatives in terms of the surface function $\eta(u)$. Equation (8) becomes approximately:

$$\chi' = -\frac{g}{c}\eta + \frac{h^2}{2}\chi''' - \frac{1}{2c}(\chi')^2 \approx -\frac{g}{c}\eta - \frac{h^2g}{2c}\eta'' - \frac{g^2}{2c^3}\eta^2. \quad (14)$$

Using similar approximations, we can eliminate $\chi'(u)$ and its higher derivatives from the surface equation (13):

$$(h + \eta) \left(-\frac{g}{c}\eta - \frac{h^2g}{2c}\eta'' - \frac{g^2}{2c^3}\eta^2 \right) + \frac{h^3g}{6c}\eta'' + c\eta = 0, \quad (15)$$

where some terms involving non-linearity of higher than 2 or involving higher order derivatives have been discarded. Collecting the leading terms, we obtain:

$$\left(1 - \frac{gh}{c^2} \right) \eta - \frac{gh^3}{3c^2}\eta'' - \frac{g}{c^2} \left(1 + \frac{gh}{2c^2} \right) \eta^2 = 0. \quad (16)$$

For the second two terms, *Fetter and Walecka* argue that it is consistent to approximate $gh \approx c^2$, which reduces (16) to

$$\left(1 - \frac{hg}{c^2} \right) \eta(u) - \frac{h^2}{3}\eta''(u) - \frac{3}{2h} [\eta(u)]^2 = 0. \quad (17)$$

Your text shows that a solution to Eq. (17) (corresponding to Eq. 56.30 of the text), with the initial condition $\eta(0) = \eta_0$ and $\eta'(0) = 0$, is the solitary wave form:

$$\zeta(x, t) = \eta(x - ct) = \eta_0 \operatorname{sech}^2 \left(\sqrt{\frac{3\eta_0}{h}} \frac{x - ct}{2h} \right), \quad (18)$$

with

$$c = \sqrt{\frac{gh}{1 - \eta_0/h}} \approx \sqrt{gh} \left(1 + \frac{\eta_0}{2h} \right). \quad (19)$$

The “standard” form of the related Korteweg-de Vries equation[2] is given in terms of the scaled variables \bar{t} and \bar{x} in terms of the function $\eta(\bar{x}, \bar{t})$ by

$$\frac{\partial \eta}{\partial \bar{t}} + 6\eta \frac{\partial \eta}{\partial \bar{x}} + \frac{\partial^3 \eta}{\partial \bar{x}^3} = 0, \quad (20)$$

which has a solution

$$\eta(\bar{x}, \bar{t}) = \frac{\beta}{2} \operatorname{sech}^2 \left[\frac{\sqrt{\beta}}{2} (\bar{x} - \beta \bar{t}) \right]. \quad (21)$$

This form is related to our results in the following way.

$$\beta = 2\eta_0, \quad \bar{x} = \sqrt{\frac{3}{2h}} \frac{x}{h}, \quad \text{and} \quad \bar{t} = \sqrt{\frac{3}{2h}} \frac{ct}{2\eta_0 h}. \quad (22)$$

To show how the reduced equation (17) is related to the Korteweg-de Vries equation, we first take the u derivative to find:

$$\frac{\eta_0}{h} \eta' - \frac{h^2}{3} \eta''' - \frac{3}{h} \eta \eta' = 0, \quad (23)$$

where we have used the relation

$$\frac{\eta_0}{h} = 1 - \frac{gh}{c^2}. \quad (24)$$

Then we notice that

$$\frac{\partial \eta}{\partial t} = -c \frac{d\eta}{du} \quad \text{and} \quad \frac{\partial \eta}{\partial x} = \frac{d\eta}{du}, \quad (25)$$

so that Eq. (23) can be written:

$$-\frac{\eta_0}{ch} \frac{\partial \eta}{\partial t} - \frac{h^2}{3} \frac{\partial^3 \eta}{\partial x^3} - \frac{3}{h} \eta \frac{\partial \eta}{\partial x} = 0. \quad (26)$$

Substituting the transformation (22) into this partial differential equation yields the Korteweg-de Vries equation (20).

References

- [1] Alexander L. Fetter and John Dirk Walecka, **Theoretical Mechanics of Particles and Continua**, (McGraw Hill, 1980), Chapt. 10.
- [2] Websites concerning solitons:
 - <http://www.ma.hw.ac.uk/solitons/>,
 - <http://www.usf.uni-osnabrueck.de/~kbrauer/solitons.html>,
 - <http://www.math.h.kyoto-u.ac.jp/~takasaki/soliton-lab/gallery/index-e.html>