

**PHY 711 Classical Mechanics and  
Mathematical Methods  
10-10:50 AM MWF Olin 103**

**Plan for Lecture 5:**

**1. Chapter 2 – Physics described  
in an accelerated coordinate  
frame**

9/7/2012

PHY 711 Fall 2012 -- Lecture 5

1

---

---

---

---

---

---

---

---

---

---

**PHY 711 Classical Mechanics and Mathematical Methods**

MWF 10 AM-10:50 PM OPL 103 <http://www.wfu.edu/~natalie/f12phy711/>

Instructor: [Natalie Holzwarth](mailto:natalie@wfu.edu) Phone:758-5510 Office:300 OPL e-mail:[natalie@wfu.edu](mailto:natalie@wfu.edu)

**Course schedule**

(Preliminary schedule -- subject to frequent adjustment.)

Date	F&W Reading	Topic	Assignment
1 Wed, 8/29/2012	Chap. 1	Review of basic principles; Scattering theory	#1
2 Fri, 8/31/2012	Chap. 1	Scattering theory continued	#2
3 Mon, 9/03/2012	Chap. 1	Scattering theory continued	#3
4 Wed, 9/05/2012	Chap. 1 & 2	Scattering theory/Accelerated coordinate frame	#4
5 Fri, 9/07/2012	Chap. 2	Accelerated coordinate frame	#5

9/7/2012

PHY 711 Fall 2012 -- Lecture 5

2

---

---

---

---

---

---

---

---

---

---

**Special seminar today: 11-11:50 AM Olin 103**



**GROWTH OF SILICENE**

Professors Hamid Oughaddou<sup>1,2</sup> and Abdelkader Kara<sup>3</sup>

<sup>1</sup>Institut des Sciences Moléculaires d'Orsay, ISMO-CNRS, Université Paris-Sud, 91405 Orsay-France

<sup>2</sup>Département de physique, Université de Cergy-Pontoise, 95000 Cergy-Pontoise, France

<sup>3</sup>Department of Physics, University of Central Florida, Orlando, FL 3216, USA

9/7/2012

PHY 711 Fall 2012 -- Lecture 5

3

---

---

---

---

---

---

---

---

---

---

Application of Newton's laws in a coordinate system which has an angular velocity  $\omega$  and linear acceleration  $\left(\frac{d^2\mathbf{a}}{dt^2}\right)_{inertial}$

Newton's laws; Let  $\mathbf{r}$  denote the position of particle of mass  $m$ :

$$m\left(\frac{d^2\mathbf{r}}{dt^2}\right)_{inertial} = \mathbf{F}_{ext}$$

$$m\left(\frac{d^2\mathbf{r}}{dt^2}\right)_{body} = \mathbf{F}_{ext} - m\left(\frac{d^2\mathbf{a}}{dt^2}\right)_{inertial} - 2m\omega \times \left(\frac{d\mathbf{r}}{dt}\right)_{body} - m\frac{d\omega}{dt} \times \mathbf{r} - m\omega \times \omega \times \mathbf{r}$$

↑ Coriolis force      ↑ Centrifugal force

9/7/2012      PHY 711 Fall 2012 -- Lecture 5      4

---

---

---

---

---

---


---

---

---

---

Motion on the surface of the Earth:



$$\omega = \frac{2\pi}{\tau} \approx 7.3 \times 10^{-5} \text{ rad/s}$$

$$\mathbf{F}_{ext} = -\frac{GM_e m}{r^2} \hat{\mathbf{r}} + \mathbf{F}'$$

Main contributions:

$$m\left(\frac{d^2\mathbf{r}}{dt^2}\right)_{earth} = -\frac{GM_e m}{r^2} \hat{\mathbf{r}} + \mathbf{F}' - 2m\omega \times \left(\frac{d\mathbf{r}}{dt}\right)_{earth} - m\frac{d\omega}{dt} \times \mathbf{r} - m\omega \times \omega \times \mathbf{r}$$

9/7/2012      PHY 711 Fall 2012 -- Lecture 5      5

---

---

---

---

---

---

---

---

---

---

Non-inertial effects on effective gravitational "constant"

$$m\left(\frac{d^2\mathbf{r}}{dt^2}\right)_{earth} = -\frac{GM_e m}{r^2} \hat{\mathbf{r}} + \mathbf{F}' - 2m\omega \times \left(\frac{d\mathbf{r}}{dt}\right)_{earth} - m\frac{d\omega}{dt} \times \mathbf{r} - m\omega \times \omega \times \mathbf{r}$$

For  $\left(\frac{d\mathbf{r}}{dt}\right)_{earth} = 0$  and  $\left(\frac{d^2\mathbf{r}}{dt^2}\right)_{earth} = 0$ ,

$$0 = -\frac{GM_e m}{r^2} \hat{\mathbf{r}} + \mathbf{F}' - m\omega \times \omega \times \mathbf{r}$$

$$\mathbf{F}' = -m\mathbf{g}$$

$$\Rightarrow \mathbf{g} = -\frac{GM_e}{r^2} \hat{\mathbf{r}} - \omega \times \omega \times \mathbf{r} \Big|_{r=R_e}$$

$$= \left( -\frac{GM_e}{R_e^2} + \omega^2 R_e \sin^2 \theta \right) \hat{\mathbf{r}} + \sin \theta \cos \theta \omega^2 R_e \hat{\boldsymbol{\theta}}$$

↑ 9.80 m/s<sup>2</sup>      ↑ 0.03 m/s<sup>2</sup>

9/7/2012      PHY 711 Fall 2012 -- Lecture 5      6

---

---

---

---

---

---

---

---

---

---

Foucault pendulum [http://www.si.edu/Encyclopedia\\_SI/nmah/pendulum.htm](http://www.si.edu/Encyclopedia_SI/nmah/pendulum.htm)



The Foucault pendulum was displayed for many years in the Smithsonian's National Museum of American History. It is named for the French physicist Jean Foucault who first used it in 1851 to demonstrate the rotation of the earth.

9/7/2012

PHY 711 Fall 2012 -- Lecture 5

7

---

---

---

---

---

---

---


---

---

---

Equation of motion on Earth's surface

$$m \left( \frac{d^2 \mathbf{r}}{dt^2} \right)_{earth} = -\frac{GM_e m}{r^2} \hat{\mathbf{r}} + \mathbf{F}' - 2m\boldsymbol{\omega} \times \left( \frac{d\mathbf{r}}{dt} \right)_{earth} - m \frac{d\boldsymbol{\omega}}{dt} \times \mathbf{r} - m\boldsymbol{\omega} \times \boldsymbol{\omega} \times \mathbf{r}$$



9/7/2012

PHY 711 Fall 2012 -- Lecture 5

8

---

---

---

---

---

---

---

---

---

---

Foucault pendulum continued – keeping leading terms:

$$m \left( \frac{d^2 \mathbf{r}}{dt^2} \right)_{earth} \approx -\frac{GM_e m}{R_e^2} \hat{\mathbf{r}} + \mathbf{F}' - 2m\boldsymbol{\omega} \times \left( \frac{d\mathbf{r}}{dt} \right)_{earth}$$

$$-\frac{GM_e m}{r^2} \hat{\mathbf{r}} \approx -mg\hat{\mathbf{z}}$$

$$\mathbf{F}' \approx -T \sin \psi \cos \phi \hat{\mathbf{x}} - T \sin \psi \sin \phi \hat{\mathbf{y}} + T \cos \psi \hat{\mathbf{z}}$$

$$\boldsymbol{\omega} \approx -\omega \sin \theta \hat{\mathbf{x}} + \omega \cos \theta \hat{\mathbf{z}}$$

$$\boldsymbol{\omega} \times \left( \frac{d\mathbf{r}}{dt} \right)_{earth} \approx \omega (-\dot{y} \cos \theta \hat{\mathbf{x}} + (\dot{x} \cos \theta + \dot{z} \sin \theta) \hat{\mathbf{y}} - \dot{y} \cos \theta \hat{\mathbf{z}})$$

9/7/2012

PHY 711 Fall 2012 -- Lecture 5

9

---

---

---

---

---

---

---

---

---

---

Foucault pendulum continued – keeping leading terms:

$$m \left( \frac{d^2 \mathbf{r}}{dt^2} \right)_{\text{earth}} \approx -\frac{GM_{\oplus} m}{R_{\oplus}^2} \hat{\mathbf{r}} + \mathbf{F}' - 2m\boldsymbol{\omega} \times \left( \frac{d\mathbf{r}}{dt} \right)_{\text{earth}}$$

$$\begin{cases} m\ddot{x} \approx -T \sin \psi \cos \phi + 2m\omega \dot{y} \cos \theta \\ m\ddot{y} \approx -T \sin \psi \sin \phi - 2m\omega(\dot{x} \cos \theta + \dot{z} \sin \theta) \\ m\ddot{z} \approx T \cos \psi - mg + 2m\omega \dot{y} \cos \theta \end{cases}$$

Further approximation :  
 $\psi \ll 1; \quad \dot{z} \approx 0; \quad T \approx mg$   
 $m\ddot{x} \approx -mg \sin \psi \cos \phi + 2m\omega \dot{y} \cos \theta$   
 $m\ddot{y} \approx -mg \sin \psi \sin \phi - 2m\omega \dot{x} \cos \theta$

Also note that :  
 $x \approx \ell \sin \psi \cos \phi$   
 $y \approx \ell \sin \psi \sin \phi$

9/7/2012 PHY 711 Fall 2012 – Lecture 5 10

---

---

---

---

---

---

---

---

---

---

Foucault pendulum continued – coupled equations:

$$\begin{cases} \ddot{x} \approx -\frac{g}{\ell} x + 2\omega \cos \theta \dot{y} \\ \ddot{y} \approx -\frac{g}{\ell} y - 2\omega \cos \theta \dot{x} \end{cases}$$

Try to find a solution of the form :  
 $x(t) = X e^{-iqt} \quad y(t) = Y e^{-iqt}$   
 Denote  $\omega_{\perp} \equiv \omega \cos \theta$   

$$\begin{pmatrix} -q^2 + \frac{g}{\ell} & i2\omega_{\perp} q \\ -i2\omega_{\perp} q & -q^2 + \frac{g}{\ell} \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix} = 0$$

Non-trivial solutions :  
 $q_{\pm} = \alpha \pm \beta \equiv \omega_{\perp} \pm \sqrt{\omega_{\perp}^2 + \frac{g}{\ell}}$

9/7/2012 PHY 711 Fall 2012 – Lecture 5 11

---

---

---

---

---

---

---

---

---

---

Foucault pendulum continued – coupled equations:

Solution continued :  
 $x(t) = X e^{-iqt} \quad y(t) = Y e^{-iqt}$   

$$\begin{pmatrix} -q^2 + \frac{g}{\ell} & i2\omega_{\perp} q \\ -i2\omega_{\perp} q & -q^2 + \frac{g}{\ell} \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix} = 0$$

Non-trivial solutions :  
 $q_{\pm} = \alpha \pm \beta \equiv \omega_{\perp} \pm \sqrt{\omega_{\perp}^2 + \frac{g}{\ell}}$   
 Amplitude relationship :  $X = iY$

General solution with complex amplitudes C and D :  
 $x(t) = \text{Re} \{ C e^{-i(\alpha+\beta)t} + i D e^{-i(\alpha-\beta)t} \}$   
 $y(t) = \text{Re} \{ C e^{-i(\alpha+\beta)t} + D e^{-i(\alpha-\beta)t} \}$

9/7/2012 PHY 711 Fall 2012 – Lecture 5 12

---

---

---

---

---

---

---

---

---

---

General solution with complex amplitudes  $C$  and  $D$ :

$$x(t) = \operatorname{Re}\{C e^{-(\alpha+\beta)t} + i D e^{-i(\alpha-\beta)t}\}$$

$$y(t) = \operatorname{Re}\{C e^{-i(\alpha+\beta)t} + D e^{-i(\alpha-\beta)t}\}$$

$$q_{\pm} = \alpha \pm \beta \equiv \omega_{\pm} \pm \sqrt{\omega_{\pm}^2 + \frac{\xi}{\tau}}$$

$$\omega_{\pm} \approx 7 \times 10^{-3} \cos \theta \text{ rad/s} \ll \sqrt{\frac{\xi}{\tau}}$$

9/7/2012

PHY 711 Fall 2012 -- Lecture 5

13

---



---



---



---



---



---



---



---