

PHY 712 Electrodynamics
12-12:50 AM MWF via video link:

<https://wakeforest-university.zoom.us/my/natalie.holzwarth>

Plan for Lecture 25:

Continue reading Chap. 11 –

Theory of Special Relativity

- A. Lorentz transformation relations**
- B. Electromagnetic field transformations**
- C. Connection to Liénard-Wiechert potentials for constant velocity sources**

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

1

In this lecture, we will continue our discussion of Special Relativity. In particular, we will discuss how the E and B fields transform between two relatively moving reference frame. Using a particular example, we will be able to show that our results for transformed fields are consistent with the results we obtain using the analysis using the Lienard-Wiechert potentials discussed earlier.

21	Mon: 03/23/2020	Chap. 9	Radiation from localized oscillating sources	#17	03/25/2020
22	Wed: 03/25/2020	Chap. 9	Radiation from oscillating sources	#18	03/27/2020
23	Fri: 03/27/2020	Chap. 9 and 10	Radiation from oscillating sources	#19	03/30/2020
24	Mon: 03/30/2020	Chap. 11	Special Theory of Relativity	#20	04/03/2020
25	Wed: 04/01/2020	Chap. 11	Special Theory of Relativity		
26	Fri: 04/03/2020	Chap. 11	Special Theory of Relativity		
27	Mon: 04/06/2020	Chap. 14	Radiation from accelerating charged particles		
28	Wed: 04/08/2020	Chap. 14	Synchrotron radiation		
	Fri: 04/10/2020	No class	<i>Good Friday</i>		
29	Mon: 04/13/2020	Chap. 14	Synchrotron radiation		
30	Wed: 04/15/2020	Chap. 15	Radiation from collisions of charged particles		
31	Fri: 04/17/2020	Chap. 13	Cherenkov radiation		
32	Mon: 04/20/2020		Special topic: E & M aspects of superconductivity		
33	Wed: 04/22/2020		Special topic: Aspects of optical properties of materials		
34	Fri: 04/24/2020				
35	Mon: 04/27/2020				
36	Wed: 04/29/2020		Review		

04/01/2020 PHY 712 Spring 2020 -- Lecture 25 2

The homework #20 assigned last lecture is due on Friday. No new homework has been assigned.

TODAY

Online colloquium scheduled for Wednesday, April 1, 2020 --
<https://www.physics.wfu.edu/events/colloquium-microstructure-control-in-organic-and-hybrid-semiconductors-and-its-impact-on-device-performance>

Online Colloquium: “Microstructure Control in Organic and Hybrid Semiconductors and its Impact on Device Performance “

Public talk for Ph. D. defense

Mr. Andrew Zeidell, Graduate Student

Mentor: Professor Oana Jurchescu

Department of Physics

Wake Forest University

Wednesday, April 1, 2020 at 3:00 PM

Video conference link: (available starting at 2:50 PM)

<https://wakeforest-university.zoom.us/j/534312421>

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

3


Please remember to attend the online lecture by Andrew Zeidell who will be describing his Ph. D. thesis work. Please notice the zoom link mentioned above and on the colloquium announcement.

Lorentz transformations

Convenient notation :

$$\beta_v \equiv \frac{v}{c}$$

$$\gamma_v \equiv \frac{1}{\sqrt{1 - \beta_v^2}}$$



Stationary frame

Moving frame

ct	=	$\gamma(ct' + \beta x')$
x	=	$\gamma(x' + \beta ct')$
y	=	y'
z	=	z'

We will continue to use the stationary and moving reference frames introduced in the previous lecture. In this case, the relative motion is along the x -axis. Of course, there is nothing special about this choice, but we will use it throughout this lecture.

Lorentz transformations -- continued

For the moving frame with $\mathbf{v} = v\hat{\mathbf{x}}$:

$$\mathcal{L}_v = \begin{pmatrix} \gamma_v & \gamma_v\beta_v & 0 & 0 \\ \gamma_v\beta_v & \gamma_v & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad \mathcal{L}_v^{-1} = \begin{pmatrix} \gamma_v & -\gamma_v\beta_v & 0 & 0 \\ -\gamma_v\beta_v & \gamma_v & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} ct \\ x \\ y \\ z \end{pmatrix} = \mathcal{L}_v \begin{pmatrix} ct' \\ x' \\ y' \\ z' \end{pmatrix} \quad \begin{pmatrix} ct' \\ x' \\ y' \\ z' \end{pmatrix} = \mathcal{L}_v^{-1} \begin{pmatrix} ct \\ x \\ y \\ z \end{pmatrix}$$

Notice :

$$c^2t^2 - x^2 - y^2 - z^2 = c^2t'^2 - x'^2 - y'^2 - z'^2$$

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

5

This slide reviews the transformations of the time and position 4-vector.

Special theory of relativity and Maxwell's equations

Continuity equation: $\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{J} = 0$

Lorenz gauge condition: $\frac{1}{c} \frac{\partial \Phi}{\partial t} + \nabla \cdot \mathbf{A} = 0$

Potential equations: $\frac{1}{c^2} \frac{\partial^2 \Phi}{\partial t^2} - \nabla^2 \Phi = 4\pi\rho$

$$\frac{1}{c^2} \frac{\partial^2 \mathbf{A}}{\partial t^2} - \nabla^2 \mathbf{A} = \frac{4\pi}{c} \mathbf{J}$$

Field relations: $\mathbf{E} = -\nabla\Phi - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t}$

$$\mathbf{B} = \nabla \times \mathbf{A}$$

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

6

This slide reviews the relevant equations for the continuity of our sources, and for Maxwell's equations in terms of the scalar and vector potentials, and for the relationship of the E and B fields to the scalar and vector potentials.

More 4-vectors:

$$\alpha = \{0, 1, 2, 3\}$$

Time and position :

$$\begin{pmatrix} ct \\ x \\ y \\ z \end{pmatrix} \Rightarrow x^\alpha$$

Charge and current :

$$\begin{pmatrix} c\rho \\ J_x \\ J_y \\ J_z \end{pmatrix} \Rightarrow J^\alpha$$

Vector and scalar potentials :

$$\begin{pmatrix} \Phi \\ A_x \\ A_y \\ A_z \end{pmatrix} \Rightarrow A^\alpha$$

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

7

Here we identify 4-vectors of time-position, charge and current sources, and scalar and vector potentials.

Lorentz transformations

$$\mathcal{L}_v = \begin{pmatrix} \gamma_v & \gamma_v \beta_v & 0 & 0 \\ \gamma_v \beta_v & \gamma_v & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Time and space :

$$x^\alpha = \mathcal{L}_v x'^\alpha \equiv \mathcal{L}_v^{\alpha\beta} x'^\beta$$

Charge and current :

$$J^\alpha = \mathcal{L}_v J'^\alpha \equiv \mathcal{L}_v^{\alpha\beta} J'^\beta$$

Vector and scalar potential : $A^\alpha = \mathcal{L}_v A'^\alpha \equiv \mathcal{L}_v^{\alpha\beta} A'^\beta$



Repeated index
summation
convention

It is reasonable to postulate that each of these three 4-vectors transform from one reference frame to another with the Lorentz transformation.

4-vector relationships

$$\begin{pmatrix} ct \\ x \\ y \\ z \end{pmatrix} \Leftrightarrow \begin{pmatrix} A^0 \\ A^1 \\ A^2 \\ A^3 \end{pmatrix} \Leftrightarrow (A^0, \mathbf{A}): \text{ upper index 4 - vector } A^\alpha \text{ for } (\alpha = 0, 1, 2, 3)$$

Keeping track of signs -- lower index 4 - vector $A_\alpha = (A^0, -\mathbf{A})$

Derivative operators (defined with different sign convention):

$$\partial^\alpha = \left(\frac{\partial}{c\partial t}, -\nabla \right) \quad \partial_\alpha = \left(\frac{\partial}{c\partial t}, \nabla \right)$$

In addition to the 4-vectors we have defined up to now, which are written with an upper index alpha, we will also need to define a lower index version of the 4-vector which just means that the space part is taken with a minus sign. We also need a notation for derivatives with respect to time and space given with the partial symbol. It turns out that for consistency, the upper and lower signs needed for the derivative operator, the upper and lower signs must be given as indicated. While Jackson's conventions are consistent throughout his text, other textbooks may use other sign conventions.

Special theory of relativity and Maxwell's equations

Continuity equation: $\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{J} = 0 \quad \rightarrow \quad \partial_\alpha J^\alpha = 0$

Lorenz gauge condition: $\frac{1}{c} \frac{\partial \Phi}{\partial t} + \nabla \cdot \mathbf{A} = 0 \quad \rightarrow \quad \partial_\alpha A^\alpha = 0$

Potential equations: $\left. \begin{aligned} \frac{1}{c^2} \frac{\partial^2 \Phi}{\partial t^2} - \nabla^2 \Phi &= 4\pi \rho \\ \frac{1}{c^2} \frac{\partial^2 \mathbf{A}}{\partial t^2} - \nabla^2 \mathbf{A} &= \frac{4\pi}{c} \mathbf{J} \end{aligned} \right\} \partial_\alpha \partial^\alpha A^\beta = \frac{4\pi}{c} J^\beta$

Field relations: $\mathbf{E} = -\nabla \Phi - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t} \quad \rightarrow ??$
 $\mathbf{B} = \nabla \times \mathbf{A}$

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

10

Here we exercise our new notation to write the important equations. I have to admit the new notation looks quite compact, (pretty, intriguing?) But what about the E and B fields, how does the new notation work for them?

Electric and Magnetic field relationships

$$\mathbf{E} = -\nabla\Phi - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t}$$

$$E_x = -\frac{\partial\Phi}{\partial x} - \frac{\partial A_x}{c\partial t} = -(\partial^0 A^1 - \partial^1 A^0)$$

$$E_y = -\frac{\partial\Phi}{\partial y} - \frac{\partial A_y}{c\partial t} = -(\partial^0 A^2 - \partial^2 A^0)$$

$$E_z = -\frac{\partial\Phi}{\partial z} - \frac{\partial A_z}{c\partial t} = -(\partial^0 A^3 - \partial^3 A^0)$$

$$\mathbf{B} = \nabla \times \mathbf{A}$$

$$B_x = \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} = -(\partial^2 A^3 - \partial^3 A^2)$$

$$B_y = \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} = -(\partial^3 A^1 - \partial^1 A^3)$$

$$B_z = \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} = -(\partial^1 A^2 - \partial^2 A^1)$$

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

11

Writing out the 6 equations for all of the E and B field components, we see that the new notation has a very nice pattern, but each field component has two indices!!! We can thus conclude that the 6 E and B field components are part of a 4x4 matrix or tensor.

Field strength tensor $F^{\alpha\beta} \equiv (\partial^\alpha A^\beta - \partial^\beta A^\alpha)$

$$F^{\alpha\beta} \equiv \begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & -B_z & B_y \\ E_y & B_z & 0 & -B_x \\ E_z & -B_y & B_x & 0 \end{pmatrix} \quad F'^{\alpha\beta} \equiv \begin{pmatrix} 0 & -E'_x & -E'_y & -E'_z \\ E'_x & 0 & -B'_z & B'_y \\ E'_y & B'_z & 0 & -B'_x \\ E'_z & -B'_y & B'_x & 0 \end{pmatrix}$$

Transformation of field strength tensor

$$F^{\alpha\beta} = \mathcal{L}_v^{\alpha\gamma} F'^{\gamma\delta} \mathcal{L}_v^{\delta\beta} \quad \mathcal{L}_v = \begin{pmatrix} \gamma_v & \gamma_v \beta_v & 0 & 0 \\ \gamma_v \beta_v & \gamma_v & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$F^{\alpha\beta} = \begin{pmatrix} 0 & -E'_x & -\gamma_v(E'_y + \beta_v B'_z) & -\gamma_v(E'_z - \beta_v B'_y) \\ E'_x & 0 & -\gamma_v(B'_z + \beta_v E'_y) & \gamma_v(B'_y - \beta_v E'_z) \\ \gamma_v(E'_y + \beta_v B'_z) & \gamma_v(B'_z + \beta_v E'_y) & 0 & -B'_x \\ \gamma_v(E'_z - \beta_v B'_y) & -\gamma_v(B'_y - \beta_v E'_z) & B'_x & 0 \end{pmatrix}$$

04/01/2020 PHY 712 Spring 2020 -- Lecture 25 12

Therefore we can define the field strength tensor and assign each of the 6 field components and their negative values to an entry in the 4x4 field strength tensor. From this logic, we can then deduce that the field strength tensor transforms as a tensor with a Lorentz transformation sandwich. Evaluating the multiplication of the three matrices, we obtain the result given on the last line. This is related to your homework problem due Friday.

Inverse transformation of field strength tensor

$$F'^{\alpha\beta} = \mathcal{L}_v^{-1\alpha\gamma} F^{\gamma\delta} \mathcal{L}_v^{-1\delta\beta}$$

$$\mathcal{L}_v^{-1} = \begin{pmatrix} \gamma_v & -\gamma_v\beta_v & 0 & 0 \\ -\gamma_v\beta_v & \gamma_v & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$F'^{\alpha\beta} = \begin{pmatrix} 0 & -E_x & -\gamma_v(E_y - \beta_v B_z) & -\gamma_v(E_z + \beta_v B_y) \\ E_x & 0 & -\gamma_v(B_z - \beta_v E_y) & \gamma_v(B_y + \beta_v E_z) \\ \gamma_v(E_y - \beta_v B_z) & \gamma_v(B_z - \beta_v E_y) & 0 & -B_x \\ \gamma_v(E_z + \beta_v B_y) & -\gamma_v(B_y + \beta_v E_z) & B_x & 0 \end{pmatrix}$$

Summary of results:

$$E'_x = E_x$$

$$B'_x = B_x$$

$$E'_y = \gamma_v(E_y - \beta_v B_z)$$

$$B'_y = \gamma_v(B_y + \beta_v E_z)$$

$$E'_z = \gamma_v(E_z + \beta_v B_y)$$

$$B'_z = \gamma_v(B_z - \beta_v E_y)$$

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

13

Using the same logic, it is possible to evaluate the inverse transformation. The last result is the same as given in Jackson Eq. 11.148.

Comparison of the two transformations

$$F'^{\alpha\beta} = \mathcal{L}'^{\alpha\gamma} F^{\gamma\delta} \mathcal{L}'^{\delta\beta}$$

$$\mathcal{L}' = \begin{pmatrix} \gamma_v & \gamma_v \beta_v & 0 & 0 \\ \gamma_v \beta_v & \gamma_v & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$F'^{\alpha\beta} = \begin{pmatrix} 0 & -E'_x & -\gamma_v(E'_y + \beta_v B'_z) & -\gamma_v(E'_z - \beta_v B'_y) \\ E'_x & 0 & -\gamma_v(B'_z + \beta_v E'_y) & \gamma_v(B'_y - \beta_v E'_z) \\ \gamma_v(E'_y + \beta_v B'_z) & \gamma_v(B'_z + \beta_v E'_y) & 0 & -B'_x \\ \gamma_v(E'_z - \beta_v B'_y) & -\gamma_v(B'_y - \beta_v E'_z) & B'_x & 0 \end{pmatrix}$$

$$F'^{\alpha\beta} = \mathcal{L}'^{-1\alpha\gamma} F^{\gamma\delta} \mathcal{L}'^{\delta\beta}$$

$$\mathcal{L}'^{-1} = \begin{pmatrix} \gamma_v & -\gamma_v \beta_v & 0 & 0 \\ -\gamma_v \beta_v & \gamma_v & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$F'^{\alpha\beta} = \begin{pmatrix} 0 & -E_x & -\gamma_v(E_y - \beta_v B_z) & -\gamma_v(E_z + \beta_v B_y) \\ E_x & 0 & -\gamma_v(B_z - \beta_v E_y) & \gamma_v(B_y + \beta_v E_z) \\ \gamma_v(E_y - \beta_v B_z) & \gamma_v(B_z - \beta_v E_y) & 0 & -B_x \\ \gamma_v(E_z + \beta_v B_y) & -\gamma_v(B_y + \beta_v E_z) & B_x & 0 \end{pmatrix}$$

04/01/2020

PHY 712 Spring 2020 -- Lecture 25


14

Comparing the various transformations.

Example:

Fields in moving frame :

$$\mathbf{E}' = \frac{q}{r'^3} (x' \hat{\mathbf{x}} + y' \hat{\mathbf{y}}) = \frac{q(-vt' \hat{\mathbf{x}} + b \hat{\mathbf{y}})}{((-vt')^2 + b^2)^{3/2}}$$

$$\mathbf{B}' = 0$$


Fields in stationary frame :

$$E_x = E'_x \qquad B_x = B'_x$$


$$E_y = \gamma_v (E'_y + \beta_v B'_z) \qquad B_y = \gamma_v (B'_y - \beta_v E'_z)$$

$$E_z = \gamma_v (E'_z - \beta_v B'_y) \qquad B_z = \gamma_v (B'_z + \beta_v E'_y)$$

04/01/2020 PHY 712 Spring 2020 -- Lecture 25 15

Now, consider a particular example discussed in Section 11.10 of Jackson. A particle sits at the origin of the moving frame. The E and B fields are measured at the point b yhat in the stationary frame. What are the values of the fields measured in the stationary frame?

Example:



Fields in moving frame :

$$\mathbf{E}' = \frac{q}{r'^3} (x' \hat{\mathbf{x}} + y' \hat{\mathbf{y}}) = \frac{q(-vt' \hat{\mathbf{x}} + b \hat{\mathbf{y}})}{\left((-vt')^2 + b^2\right)^{3/2}}$$

$$\mathbf{B}' = 0$$

Fields in stationary frame :

$$E_x = E'_x = \frac{q(-vt')}{\left((-vt')^2 + b^2\right)^{3/2}}$$


$$E_y = \gamma_v (E'_y) = \frac{q(\gamma_v b)}{\left((-vt')^2 + b^2\right)^{3/2}}$$

$$B_z = \gamma_v (\beta_v E'_y) = \frac{q(\gamma_v \beta_v b)}{\left((-vt')^2 + b^2\right)^{3/2}}$$

04/01/2020 PHY 712 Spring 2020 -- Lecture 25 16

It is easy to write the fields in the moving frame, since the particle is stationary in that frame. Then we use the transformation equations to find the fields in the stationary frame. We are not quite done, because the expressions involve the time measured in the moving frame.

Example:



Fields in moving frame :

$$\mathbf{E}' = \frac{q}{r'^3} (x' \hat{\mathbf{x}} + y' \hat{\mathbf{y}}) = \frac{q(-vt' \hat{\mathbf{x}} + b \hat{\mathbf{y}})}{((-vt')^2 + b^2)^{3/2}}$$

$$\mathbf{B}' = 0$$

Fields in stationary frame :

$$E_x = E'_x = \frac{q(-v\gamma_v t)}{((-v\gamma_v t)^2 + b^2)^{3/2}}$$

$$E_y = \gamma_v (E'_y) = \frac{q(\gamma_v b)}{((-v\gamma_v t)^2 + b^2)^{3/2}}$$

$$B_z = \gamma_v (\beta_v E'_y) = \frac{q(\gamma_v \beta_v b)}{((-v\gamma_v t)^2 + b^2)^{3/2}}$$

Expression in terms of consistent coordinates

$$t' = \gamma_v t$$


04/01/2020

PHY 712 Spring 2020 -- Lecture 25

17

Using the time-coordinate transformation we can then write the fields measured in the stationary frame in terms of the time appropriate to that frame.

$$E_y = \frac{q(\gamma_v b)}{\left((-v\gamma_v t)^2 + b^2\right)^{3/2}} = \frac{q(\gamma_v b)}{\left((\gamma_v^2 - 1)c^2 t^2 + b^2\right)^{3/2}}$$



04/01/2020

PHY 712 Spring 2020 -- Lecture 25

18

This plot shows the y component of the electric field as measured in the stationary frame plotted as a function of time. For large gamma, there is a large peak at $t=0$.

Examination of this system from the viewpoint of the
the Liènard-Wiechert potentials (temporarily keeping SI units)

$$\rho(\mathbf{r}, t) = q\delta^3(\mathbf{r} - \mathbf{R}_q(t)) \quad \mathbf{J}(\mathbf{r}, t) = q\dot{\mathbf{R}}_q(t)\delta^3(\mathbf{r} - \mathbf{R}_q(t)) \quad \dot{\mathbf{R}}_q(t) = \frac{d\mathbf{R}_q(t)}{dt}$$

$$\Phi(\mathbf{r}, t) = \frac{1}{4\pi\epsilon_0} \iiint d^3r' dt' \frac{\rho(\mathbf{r}', t')}{|\mathbf{r} - \mathbf{r}'|} \delta(t' - (t - |\mathbf{r} - \mathbf{r}'|/c))$$

$$\mathbf{A}(\mathbf{r}, t) = \frac{1}{4\pi\epsilon_0 c^2} \iiint d^3r' dt' \frac{\mathbf{J}(\mathbf{r}', t')}{|\mathbf{r} - \mathbf{r}'|} \delta(t' - (t - |\mathbf{r} - \mathbf{r}'|/c))$$

Evaluating integral over t' :

$$\int_{-\infty}^{\infty} dt' f(t') \delta(t' - (t - |\mathbf{r} - \mathbf{R}_q(t')|/c)) = \frac{f(t_r)}{1 - \frac{\mathbf{R}_q(t_r) \cdot (\mathbf{r} - \mathbf{R}_q(t_r))}{c |\mathbf{r} - \mathbf{R}_q(t_r)|}},$$

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

19

Do these results make sense? In order to check the results, we can calculate the fields directly in the stationary frame using the methods we discussed several lectures ago using the Lienard-Wiechert potentials. Here we review some of those equations.

Examination of this system from the viewpoint of the the Liénard-Wiechert potentials – continued (SI units)

$$\Phi(\mathbf{r}, t) = \frac{q}{4\pi\epsilon_0} \frac{1}{R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}}$$

$$\mathbf{A}(\mathbf{r}, t) = \frac{q}{4\pi\epsilon_0 c^2} \frac{\mathbf{v}}{R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}}$$

where $\mathbf{R} = \mathbf{r} - \mathbf{R}_q(t_r)$ $\mathbf{v} = \frac{d\mathbf{R}_q(t_r)}{dt_r}$

$$\mathbf{E}(\mathbf{r}, t) = -\nabla\Phi(\mathbf{r}, t) - \frac{\partial\mathbf{A}(\mathbf{r}, t)}{\partial t}$$

$$\mathbf{B}(\mathbf{r}, t) = \nabla \times \mathbf{A}(\mathbf{r}, t)$$

More equations.

Examination of this system from the viewpoint of the
the Liénard-Wiechert potentials – continued (SI units)

$$\mathbf{E}(\mathbf{r}, t) = \frac{q}{4\pi\epsilon_0} \frac{1}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^3} \left[\left(\mathbf{R} - \frac{\mathbf{v}R}{c}\right) \left(1 - \frac{v^2}{c^2}\right) + \left(\mathbf{R} \times \left\{ \left(\mathbf{R} - \frac{\mathbf{v}R}{c}\right) \times \frac{\dot{\mathbf{v}}}{c^2} \right\}\right) \right]$$

$$\mathbf{B}(\mathbf{r}, t) = \frac{q}{4\pi\epsilon_0 c^2} \left[\frac{-\mathbf{R} \times \mathbf{v}}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^3} \left(1 - \frac{v^2}{c^2} + \frac{\dot{\mathbf{v}} \cdot \mathbf{R}}{c^2}\right) - \frac{\mathbf{R} \times \dot{\mathbf{v}} / c}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^2} \right]$$

$$\mathbf{B}(\mathbf{r}, t) = \frac{\mathbf{R} \times \mathbf{E}(\mathbf{r}, t)}{cR}.$$

Finally the E and B fields obtained from that analysis.

Examination of this system from the viewpoint of the
the Liénard-Wiechert potentials – (**Gaussian units**)

$$\mathbf{E}(\mathbf{r}, t) = \frac{q}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^3} \left[\left(\mathbf{R} - \frac{\mathbf{v}R}{c}\right) \left(1 - \frac{v^2}{c^2}\right) + \left(\mathbf{R} \times \left\{ \left(\mathbf{R} - \frac{\mathbf{v}R}{c}\right) \times \frac{\dot{\mathbf{v}}}{c^2} \right\}\right) \right]$$

$$\mathbf{B}(\mathbf{r}, t) = \frac{q}{c} \left[\frac{-\mathbf{R} \times \mathbf{v}}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^3} \left(1 - \frac{v^2}{c^2} + \frac{\dot{\mathbf{v}} \cdot \mathbf{R}}{c^2}\right) - \frac{\mathbf{R} \times \dot{\mathbf{v}} / c}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^2} \right]$$

$$\mathbf{B}(\mathbf{r}, t) = \frac{\mathbf{R} \times \mathbf{E}(\mathbf{r}, t)}{R}.$$

Here are the equations in cgs Gaussian units that we are now using.

Examination of this system from the viewpoint of the
the Liénard-Wiechert potentials – continued (**Gaussian units**)

$$\mathbf{E}(\mathbf{r}, t) = \frac{q}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^3} \left[\left(\mathbf{R} - \frac{\mathbf{v}R}{c}\right) \left(1 - \frac{v^2}{c^2}\right) \right]$$

For our example:

$$\mathbf{R}_q(t_r) = vt_r \hat{\mathbf{x}} \quad \mathbf{r} = b\hat{\mathbf{y}}$$

$$\mathbf{R} = b\hat{\mathbf{y}} - vt_r \hat{\mathbf{x}} \quad R = \sqrt{v^2 t_r^2 + b^2}$$

$$\mathbf{B}(\mathbf{r}, t) = \frac{q}{c} \left[\frac{-\mathbf{R} \times \mathbf{v}}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^3} \left(1 - \frac{v^2}{c^2}\right) \right]$$

$$\mathbf{v} = v\hat{\mathbf{x}} \quad t_r = t - \frac{R}{c}$$

This should be equivalent to the result given in Jackson (11.152):

$$\mathbf{E}(x, y, z, t) = \mathbf{E}(0, b, 0, t) = q \frac{-v\gamma t \hat{\mathbf{x}} + \gamma b \hat{\mathbf{y}}}{(b^2 + (v\gamma t)^2)^{3/2}}$$

$$\mathbf{B}(x, y, z, t) = \mathbf{B}(0, b, 0, t) = q \frac{\gamma \beta b \hat{\mathbf{z}}}{(b^2 + (v\gamma t)^2)^{3/2}}$$

04/01/2020

PHY 712 Spring 2020 -- Lecture 25

23

Now to evaluate the equations, we need to consider the constant velocity trajectory of our example. We will continue this discussion on Friday.