

# A SYSTEM OF ODES FOR A PERTURBATION OF A MINIMAL MASS SOLITON

JEREMY L. MARZUOLA, SARAH RAYNOR, AND GIDEON SIMPSON

ABSTRACT. We study soliton solutions to the nonlinear Schrödinger equation (NLS) with a saturated nonlinearity. NLS with such a nonlinearity is known to possess a minimal mass soliton. We consider a small perturbation of a minimal mass soliton and identify a system of ODEs extending the work of [20], which models the behavior of the perturbation for short times. We then provide numerical evidence that under this system of ODEs there are two possible dynamical outcomes, in accord with the conclusions of [43]. Generically, initial data which supports a soliton structure appears to oscillate, with oscillations centered on a stable soliton. For initial data which is expected to disperse, the finite dimensional dynamics initially follow the unstable portion of the soliton curve.

## 1. INTRODUCTION

The nonlinear Schrödinger equation (NLS) in  $\mathbb{R}^d \times \mathbb{R}^+$ :

$$(1.1) \quad iu_t + \Delta u + g(|u|^2)u = 0, \quad u(x, 0) = u_0(x)$$

emerges as a leading order description of important physical systems in optics, many body quantum systems, hydrodynamics, and plasma physics. Moreover, it is a canonical equation demonstrating the competition between nonlinearity and dispersion.

In many applications, the leading order approximation of the nonlinearity,  $g(s)$ , is the power nonlinearity  $g(s) = \pm s^\sigma$ . For example,  $g(s) = s$ , yields the focusing cubic NLS equation. Cubic NLS appears in many contexts, including many body quantum quantum systems, when the Hartree equation is approximated with pairwise delta function interactions. It similarly appears in nonlinear optics if the higher order corrections to the nonlinear index of refraction are neglected.

Such approximations may lead to non-physical predictions. In the case of the power-law nonlinearity, if  $\sigma d \geq 2$ , initial data with mass ( $L^2$  norm) in excess of the ground state may blow up in finite time [58]. However, experiments in the optical setting show that there is no “singularity” to speak of; the intensity of the solutions remains finite [34].

The model equation can be corrected to suppress singularity formation by replacing the power nonlinearity with a *saturated* nonlinearity. Ideally, such a nonlinearity allows potentially unstable behavior at low intensity but regularizes it at high intensity. One example of such a nonlinearity is the cubic-quintic, where  $g(s) = s - \gamma s^2$ . The cubic-quintic NLS (CQNLS) equation appears as a correction to the many body system that

includes (repulsive) three body delta function interactions [8, 9]. CQNLS also appears in models of super fluids, [33], and in nonlinear optics, [23, 45, 62].

Another saturated nonlinearity appearing optics literature is  $g(s) = s/(1+s)$ , [59]. This nonlinearity was used in [24] to study turbulence for a one dimensional, non-integrable equation without singularity formation. In laser plasma interactions,  $g(s) = 1 - e^{-s}$  is used, [31]. In all cases, the Taylor expansion of the nonlinearity leads to CQNLS at second order.

In this work we consider saturated nonlinearities of the form

$$(1.2) \quad g(s) = s^{\frac{q}{2}} \frac{s^{\frac{p-q}{2}}}{1 + s^{\frac{p-q}{2}}},$$

where  $2 + \frac{4}{d-2} > p > 2 + \frac{4}{d} > \frac{4}{d} > q > 0$  for  $d \geq 3$  and  $\infty > p > 2 + \frac{4}{d} > \frac{4}{d} > q > 0$  for  $d < 3$ . For  $|u|$  large, (1.1) behaves as though it were  $L^2$  subcritical while for  $|u|$  small, it behaves as though it were  $L^2$  supercritical. Note that this choice of saturated nonlinearity is focusing for all values of  $|u|$ . For our purposes,  $p$  must be chosen substantially larger than the  $L^2$  critical exponent,  $\frac{4}{d}$ , in order to allow sufficient regularity when linearizing the equation. For our numerical analysis, we work in one spatial dimension with the specific nonlinearity

$$(1.3) \quad g(s) = \frac{s^3}{1 + s^2}.$$

A soliton solution of (1.1) is a function  $u(t, x)$  of the form

$$(1.4) \quad u(t, x) = e^{i\omega t} \phi_\omega(x),$$

where  $\omega > 0$  and  $\phi_\omega(x)$  is a positive, spherically symmetric, exponentially decaying solution of the equation:

$$(1.5) \quad \Delta \phi_\omega - \omega \phi_\omega + g(\phi_\omega^2) \phi_\omega = 0.$$

For our particular nonlinearity, for any  $\omega > 0$  there is a unique solitary wave solution  $\phi_\omega(x)$  to (1.5), see [12] and [41].

For large  $\omega$  the solitons are stable, while for small  $\omega$  they are unstable. A precise stability criterion identifying stable and unstable regions is provided in [29] and [51], generalizing earlier work on stability in [60], [61]. This amounts to examining the relation  $\omega \mapsto \|\phi_\omega\|_{L^2}^2$ , defining a *soliton curve*. Where it is increasing as a function of  $\omega$ , the solitons are stable; where this function is decreasing the solitons are unstable. In Figure 1 we plot this curve for several common choices of  $g(s)$ .

As can be seen numerically in Figure 1, our saturated nonlinearity spawns a soliton of minimal mass. Certain asymptotic methods can be used to describe the nature of the curve—multiscale methods can provide the asymptotics as  $\omega \rightarrow 0$ , while variational methods can give the asymptotics as  $\omega \rightarrow \infty$ . However, in this work we forego an analytic description of the soliton curve and focus on the minimal mass soliton shown to exist in the numerical




FIGURE 1. Plots of the soliton curves ( $\phi(\omega)$  with respect to  $\omega$ ) for a subcritical nonlinearity, critical nonlinearity, supercritical nonlinearity, and the saturated nonlinearity (1.3). The curves for the monomial nonlinearities are found analytically, while the curves for the saturated nonlinearities are found numerically using the method discussed in Section 4.1. All are in  $d = 1$ . Here  $\omega \geq .001$ , as the supercritical and saturated cases diverge as  $\omega \rightarrow 0$ .

plot. The minimal mass solution is a distinguished case amongst the already special family of soliton solutions. Though solitons to the left and right of this minimum can readily be classified as stable or unstable, the theory fails at extrema [60,61]. This motivates a careful examination of the dynamics of the minimal mass soliton under perturbation.

In [20], Comech and Pelinovsky demonstrated that the minimal mass soliton possesses a fundamentally *nonlinear* instability. They accomplished this by finding a small perturbation that forces the solution a fixed distance away from the minimal mass soliton in finite time. Their technique reduces the problem to studying an ODE modeling the perturbation for short times. They show that the 0 solution to the ODE is unstable by appropriate choice of initial data.

The purpose of this work is to numerically explore the following conjecture:

**Conjecture 1.1.** *Though the minimal mass soliton is unstable on short time scales, we conjecture that, on longer time scales, any solutions which is initially a small perturbation of the minimal mass soliton will either disperse or eventually relax towards a nearby stable soliton.*

This conjecture is part of the more general conjecture that any solution which does not disperse as  $t \rightarrow \infty$  must eventually converge to a finite sum of stable solitons. Although the equation has a branch of unstable solitons, physically, it is expected that general solutions to (1.1) will resolve into stable solitons plus a dispersive component. This is referred to as the “soliton resolution” conjecture, a notoriously difficult problem to formulate. For nonlinearities with a specific two power structure, dynamics of this type were observed by Pelinovsky, Afanasjev, and Kivshar, who modelled the behavior of solutions near a minimal mass soliton by a second order ODE via adiabatic expansion in  $\omega$  [43]. This conjecture has also been explored numerically by Buslaev and Grikurov for two power nonlinearities in [16], where they found that a solution which is initially a perturbation of an unstable soliton tends to approach and then oscillate around a stable soliton.

By contrast, our method uses the full dynamical system of modulation parameters to find a 4-dimensional system of ODEs which is structured to allow for eventual recoupling to the continuous spectrum. Following [20], we begin with the ansatz that our solution is a small perturbation of the minimal mass soliton. We then break the perturbation into discrete and continuous spectral components relative to the linearization of the Schrödinger operator about the soliton. The discrete portion yields a four dimensional system of nonlinear ODEs. In this work, we neglect the continuous spectrum, although it may be included in a future work. We further simplify the system expanding the equations in powers of the dependent variables and dropping cubic and higher terms.

An obstacle in studying these ODEs is that the signs and magnitudes of the coefficients are not self-evident, necessitating numerical methods. We compute these values, which are intimately related to the minimal mass soliton, using the sinc spectral method. The use of the sinc function for numerically solving differential equations dates to Stenger [54]. It has been successfully used in a wide variety of linear and nonlinear, time dependent and independent, differential equations [5, 11, 14, 18, 25, 38, 46, 57]. In this work, we first numerically solve (1.5) for the soliton as a nonlinear collocation problem. We then use this information to compute the generalized kernel of the operator after linearization about the soliton.

With these coefficients in hand, we numerically integrate the ODE system, plotting the results. We find that there are two different types of behavior for the finite dimensional system, depending on the initial data. If the initial data represents a solution which our nonlinear PDE solver indicates can support a soliton, then we find that the solution is oscillatory. It is initially attracted to the stable side of the curve, and, over intermediate time scales, oscillates around a stable soliton close to the minimal mass soliton. If we initialize with the unstable conditions found in [20], the ODEs initially move in the unstable direction but quickly reverse, before commencing oscillation. On the other hand, if we begin with initial conditions which are expected to disperse as  $t \rightarrow \infty$ , the finite dimensional dynamics push the solution along the unstable soliton curve towards the value  $\omega = 0$  rather quickly. This solution agrees with the numerically computed solution of the primitive

equation (1.1) with corresponding initial data for as long as the mass conservation of the solution allows, after which our model continues to follow the unstable soliton curve but the actual solution disperses. In [43], the authors observed similar dynamics, with both oscillatory and dispersive regimes.

These ODEs are an approximation valid on a short time interval. This study is the beginning of an analysis to show that perturbations of the minimal mass soliton are attracted to the stable side of the soliton curve. In a forthcoming work we hope to show how the continuous-spectrum part of the perturbation interacts with the discrete-spectrum perturbation. Based on the work of Soffer and Weinstein, [53] we expect coupling to the continuous spectrum to cause radiation damping, which will ultimately cause the solution to have damped oscillations and select a soliton on the stable side of the curve.

This paper is organized as follows. In section 2, we introduce preliminaries and necessary definitions. In section 3, we derive the system of ODEs. In section 4, we explain our numerical methods for finding the coefficients of the ODEs. In section 5, we show the numerical solutions of the ODEs and explain our results. Finally, in section 6 we present our conclusions and plans for future work. An appendix contains details of our numerical method for computation of the soliton and related coefficients.

**ACKNOWLEDGMENTS** This project began out of a conversation with Catherine Sulem and JLM. JLM was partially funded by an NSF Postdoc at Columbia University and a Hausdorff Center Postdoc at the University of Bonn. In addition, JLM would like to thank the University of North Carolina, Chapel Hill for graciously hosting him during part of this work. SR would like to thank the University of Chicago for their hospitality while some of this work was completed. GS was funded in part by NSERC. In addition, the authors wish to thank Dmitry Pelinovsky, Mary Pugh, Catherine Sulem and Michael Weinstein for helpful comments and suggestions.

## 2. DEFINITIONS AND SETUP

Equation (1.1) is globally well-posed in  $H^1$ , the usual Sobolev space, with norm

$$\|u\|_{H^1}^2 = \|u\|_{L^2}^2 + \|\nabla u\|_{L^2}^2.$$

The global well-posedness for initial data in  $H^1$  follows from the standard well-posedness theory for semilinear Schrödinger equations. Additionally, we assume that  $u_0$  is spherically symmetric (which means even in one dimension), implying  $u(x, t) = u(|x|, t)$  for all  $t > 0$ . Proofs can be found in numerous references including [19] and [58].

For data  $u_0 \in H^1$ , there are several conserved quantities. Particularly important invariants are:

### **Conservation of Mass (or Charge):**

$$Q(u) = \frac{1}{2} \int_{\mathbb{R}^d} |u|^2 dx = \frac{1}{2} \int_{\mathbb{R}^d} |u_0|^2 dx.$$

**Conservation of Energy:**

$$E(u) = \int_{\mathbb{R}^d} |\nabla u|^2 dx - \int_{\mathbb{R}^d} G(|u|^2) dx = \int_{\mathbb{R}^d} |\nabla u_0|^2 dx - \int_{\mathbb{R}^d} G(|u_0|^2) dx,$$

where

$$G(t) = \int_0^t g(s) ds.$$

Detailed proofs of these conservation laws can be easily arrived at by using energy estimates or Noether's Theorem, which relates conservation laws to symmetries of an equation. See [58] for details.

With this type of nonlinearity, it is known that soliton solutions to NLS exist and are unique. Existence of solitary waves for nonlinearities of the type (1.2) is proved by in [12] in  $\mathbb{R}^1$  using ODE techniques and in higher dimensions by minimizing the functional

$$T(u) = \int |\nabla u|^2 dx$$

with respect to the functional

$$V(u) = \int [G(|u|^2) - \frac{\omega}{2}|u|^2] dx.$$

Then, using a minimizing sequence and Schwarz symmetrization, one infers the existence of the nonnegative, spherically symmetric, decreasing soliton solution. Once we know that minimizers are radially symmetric, uniqueness can be established via a shooting method, showing that the desired soliton occurs at only one initial value, [41].

Of great importance is the fact that  $Q_\omega := Q(\phi_\omega)$  and  $E_\omega := E(\phi_\omega)$  are differentiable with respect to  $\omega$ . This can be determined from the works of Shatah, namely [49], [50]. Differentiating (1.5),  $Q$  and  $E$  all with respect to  $\omega$ , we have the relation

$$\partial_\omega E_\omega = -\omega \partial_\omega Q_\omega.$$

Numerical computations show that if we plot  $Q_\omega$  with respect to  $\omega$  for the saturated nonlinearity, the soliton curve goes to  $\infty$  as  $\omega$  goes to 0 or  $\infty$  and has a global minimum at some  $\omega = \omega_* > 0$ ; see Figure 1.

We are interested in the stability of these explicit solutions under perturbations of the initial data. Recall the following notions of stability:

**Definition 2.1.** *The soliton is said to be **orbitally stable** if,  $\forall \epsilon > 0$ ,  $\exists \delta > 0$  such that, for any initial data  $u_0$  such that  $\|u_0 - \phi_\omega\|_{H^1} < \delta$ , for any  $t < 0$ , there is some  $\theta \in \mathbb{R}$  such that  $\|u(x, t) - e^{i\theta} \phi_\omega(x)\|_{H^1} < \epsilon$ .*

**Definition 2.2.** *The soliton is said to be **asymptotically stable**, if,  $\exists \delta > 0$  such that if  $\|u_0 - \phi_\omega\|_{H^1} < \delta$ , then  $\exists \tilde{\omega}, \tilde{\theta} > 0$  and  $\psi_0 \in H^1$  such that  $\|u(x, t) - e^{i\tilde{\omega}t + \tilde{\theta}} \phi_{\tilde{\omega}}(x) - e^{it\Delta} \psi_0\|_{H^1} \rightarrow 0$  as  $t \rightarrow \infty$ .*

**Remark 2.1.** *Orbital stability suggests that a solution to (1.1) initially near the soliton will remain near the soliton in the  $H^1$  norm. However, asymptotic stability states that the system actually selects a specific soliton to converge to as time goes to  $\infty$ . Note, technically a system can undergo a very large change in  $\omega$  and still be “asymptotically stable” provided it converges to a soliton for large time. However, asymptotic stability is usually proved using perturbation theory, which would suggest that in Definition 2.2, there would exist an  $\epsilon > 0$  depending on  $\delta$  such that*

$$|\omega - \tilde{\omega}| < \epsilon.$$

Variational techniques developed in [60], [61] and generalized to an abstract setting in [29] and [51] tell us that when  $\delta(\omega) = E_\omega + \omega Q_\omega$  is convex, or  $\delta''(\omega) > 0$ , we are guaranteed stability under small perturbations, while for  $\delta''(\omega) < 0$  we are guaranteed that the soliton is unstable under small perturbations. For brief reference on this subject, see Chapter 4 of [58]. For nonlinearities that are twice differentiable at the origin and of monomial type at infinity (which would include our saturated nonlinearities), asymptotic stability has been studied for a finite collection of strongly orbitally stable solitons by Buslaev and Perelman [17], Cuccagna [21], and Rodnianski, Soffer and Schlag [47].

At a minimum of  $Q_\omega$ , soliton instability is more subtle, because it is due solely to nonlinear effects. See [20], where this purely nonlinear instability is proved to occur by reducing the behavior of the discrete part of the spectrum to an ODE that is unstable for certain initial conditions.

**2.1. Linearization about a Soliton.** Throughout this section, we use vector notation,  $\vec{\cdot}$ , to represent complex functions. Any function written without vector notation is assumed to be real. For example, the complex valued scalar function  $u + iv$  will be written  $\begin{pmatrix} u \\ v \end{pmatrix}$ . In this notation, multiplication by  $i$  is represented by the matrix  $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ . We denote by  $\vec{\phi}_\omega$  the complex vector  $\begin{pmatrix} \phi_\omega \\ 0 \end{pmatrix}$ , where  $\phi_\omega$  is the real profile of the soliton with parameter  $\omega$ . For simplicity, we suppress the  $\omega$  subscript, writing  $\phi$  in place of  $\phi_\omega$ .

For later reference, we now explicitly characterize the linearization of NLS about a soliton solution. First consider the linear evolution of the perturbation of a soliton via the ansatz:

$$(2.1) \quad \vec{u} = e^{J\omega t}(\vec{\phi}_\omega(x) + \vec{\rho}(x, t))$$

with  $\vec{\rho} = \begin{pmatrix} \rho_{Re} \\ \rho_{Im} \end{pmatrix}$ . For the purposes of finding the linearized Hamiltonian at  $\phi_\omega$  we do not need to allow the parameters  $\theta$  and  $\omega$  to modulate, but when we develop our full system of equations in Section 3 parameter modulation will be taken into account. Inserting (2.1)

as an ansatz into (1.1) we know that since  $\phi$  is a soliton solution we have

$$(2.2) \quad J(\vec{\rho})_t + \Delta(\vec{\rho}) - \omega\vec{\rho} = -g(\phi^2)\vec{\rho} - 2g'(\phi^2)\phi^2 \begin{pmatrix} \rho_{Re} \\ 0 \end{pmatrix} + O(|\vec{\rho}|^2).$$

(This calculation is explained in more detailed at the start of Section 3.) Here we have used the following calculation of the nonlinear terms of the perturbation equation:

$$(2.3) \quad \begin{aligned} (g(|\phi + \rho|^2)(\phi + \rho) - g(|\phi|^2)\phi) &= (g(\phi^2 + 2\phi\rho_{Re} + \rho_{Re}^2 + \rho_{Im}^2)(\phi + \rho_{Re} + i\rho_{Im}) - g(\phi^2)\phi) \\ &= g'(\phi^2)(\rho_{Re}^2 + 2\phi\rho_{Re} + \rho_{Im}^2)(\phi + \rho_{Re} + i\rho_{Im}) \\ &\quad + \frac{1}{2}g''(\phi^2)(\rho_{Re}^2 + 2\phi\rho_{Re} + \rho_{Im}^2)^2(\phi + \rho_{Re} + i\rho_{Im}) + \text{h.o.t.s.} \end{aligned}$$

The linear terms will be absorbed into the linearized operator  $J\mathcal{H}_\omega$ , while the quadratic terms are handled explicitly; the  $\mathcal{O}(\rho^2)$  terms in the expansion of the equation around  $\phi_\omega$  will be denoted by  $N(\omega, \rho)$  in the sequel. In this work, after expansion in powers of  $\rho$ , we drop all terms of order greater than two.

We are interested in linearizing this equation:

$$(2.4) \quad \partial_t \begin{pmatrix} \rho_{Re} \\ \rho_{Im} \end{pmatrix} = J\mathcal{H} \begin{pmatrix} \rho_{Re} \\ \rho_{Im} \end{pmatrix} + \text{h.o.t.},$$

where

$$(2.5) \quad \mathcal{H} = \begin{pmatrix} 0 & L_- \\ -L_+ & 0 \end{pmatrix},$$

$$(2.6) \quad L_- = -\Delta + \omega - g(\phi_\omega),$$

$$(2.7) \quad L_+ = -\Delta + \omega - g(\phi_\omega) - 2g'(\phi_\omega^2)\phi_\omega^2.$$

**Definition 2.3.** *A Hamiltonian,  $\mathcal{H}$  is called admissible if the following hold:*

- 1) *There are no embedded eigenvalues in the essential spectrum,*
- 2) *The only real eigenvalue in  $[-\omega, \omega]$  is 0,*
- 3) *The values  $\pm\omega$  are non-resonant .*

**Definition 2.4.** *Let (NLS) be taken with nonlinearity  $g$ . We call  $g$  admissible if there exists a minimal mass soliton,  $\phi_{min}$ , for (NLS) and the Hamiltonian,  $\mathcal{H}$ , resulting from linearization about  $\phi_{min}$  is admissible in terms of Definition 2.3.*

The spectral properties assumed for the linearized Hamiltonian equation in order to prove stability results are typically those from Definition 2.3, see [17, 22, 48, 47, 35, 28]. However, we note that this condition is a constraint used to control the dispersive estimates necessary for perturbative analysis in analytic results, though it is sometimes possible to numerically solve this sort of problem even if Definition 2.3 does not hold; see, for example, [16].

In this work, we must simply assume that  $g$  is an admissible nonlinearity so that we must only concern ourselves with known discrete spectrum functions without coupling to



endpoint resonances or embedded eigenvalues. However, this assumption is justified by the observed dynamics. Great care must be taken in studying the spectral properties of a linearized operator; although admissibility is expected to hold generically, certain algebraic conditions on the soliton structure itself must factor into the analysis, often requiring careful numerical computations. See [22] as an introduction to such methods and the difficulties therein. To this end, in the forthcoming work [40], two of the authors will look at analytic and computational methods for verifying these spectral conditions.

**2.2. The Discrete Spectral Subspace.** We approximate perturbations of the minimal mass soliton by projecting onto the discrete spectral subspace of the linearized operator. Notationally, we refer to  $P_d$  as the projection onto the finite dimensional discrete spectral subspace  $D_\omega$  of  $H^1$  relative to  $\mathcal{H}$ . Similarly,  $P_c$  represents projection onto the continuous spectral subspace for  $\mathcal{H}$ . We now describe, in detail, the discrete spectral subspace at the minimal mass.

Let  $\omega^*$  be the value of the soliton parameter at which the minimal mass soliton occurs. It is proved in [20] (Lemma 3.8) that the discrete spectral subspace  $D_\omega$  of  $\mathcal{H}$  at  $\omega^*$  has real dimension 4. This results from the extra orthogonality at minimal mass, which gives a dimension 4 generalized kernel given by the following chain of equalities

$$\begin{aligned} L_- \phi_{\omega^*} &= 0, \\ L_+(-\phi'_{\omega^*}) &= \phi_{\omega^*}, \\ L_- \alpha &= -\phi'_{\omega^*}, \\ L_+ \beta &= \alpha. \end{aligned}$$

The functions  $\vec{e}_1 = \begin{pmatrix} 0 \\ \phi_\omega \end{pmatrix}$  and  $\vec{e}_2 = \begin{pmatrix} \phi'_\omega \\ 0 \end{pmatrix}$  are in the generalized kernel of  $\mathcal{H}$  at every  $\omega$ . Clearly,  $\vec{e}_1$  is purely imaginary and  $\vec{e}_2$  is real. In addition to  $\vec{e}_1$  and  $\vec{e}_2$ , at  $\omega^*$  there are two more linearly independent elements of  $D_\omega$ , the purely imaginary  $\vec{e}_3 = \begin{pmatrix} 0 \\ \alpha \end{pmatrix}$  and the purely real  $\vec{e}_4 = \begin{pmatrix} \beta \\ 0 \end{pmatrix}$ .

Applying [20] (Lemma 3.9),  $\vec{e}_3$  and  $\vec{e}_4$  can be extended as continuous functions of  $\omega$  in such a way that  $\vec{e}_3(\omega)$  is purely imaginary,  $\vec{e}_4(\omega)$  is purely real. We write

$$\vec{e}_3(\omega) = \begin{pmatrix} 0 \\ \alpha(\omega) \end{pmatrix}$$

and

$$\vec{e}_4(\omega) = \begin{pmatrix} \beta(\omega) \\ 0 \end{pmatrix},$$

with  $\alpha$  and  $\beta$  real-valued functions. The linearized operator, restricted to this subspace, is

$$(2.8) \quad J\mathcal{H}(\omega)|_{D_\omega} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & a(\omega) & 0 \end{pmatrix},$$

where  $a(\omega)$  is a differentiable function that is equal to 0 at  $\omega^*$ .

Before proceeding to the derivation of the ODEs, it is helpful to make a minor change of basis. Our goal is that, in the new basis,  $\{\vec{e}_1, \vec{e}_2, \vec{e}_3, \vec{e}_4\}$ ,  $\langle \vec{e}_1, \vec{e}_3 \rangle = 0$ , which will make it easier for us to compute the dual basis. Replace  $\vec{e}_3$  by

$$\begin{aligned} \vec{e}_3 &= \vec{e}_3 - \frac{\langle \vec{e}_1, \vec{e}_3 \rangle}{\|\vec{e}_1\|^2} \vec{e}_1 \\ &= \begin{bmatrix} 0 \\ \tilde{\alpha} \end{bmatrix}. \end{aligned}$$

To preserve the relationship  $\vec{e}_3 = JH_\omega \vec{e}_4$ , we need to replace  $\vec{e}_4$  by

$$\begin{aligned} \vec{e}_4 &= \vec{e}_4 - \frac{\langle \vec{e}_1, \vec{e}_3 \rangle}{\|\vec{e}_1\|^2} \vec{e}_2 \\ &= \begin{bmatrix} \tilde{\beta} \\ 0 \end{bmatrix}. \end{aligned}$$

To preserve the relationship  $JH_\omega \vec{e}_3 = \vec{e}_2 + a(\omega) \vec{e}_4$ , we replace  $\vec{e}_2$  by

$$\begin{aligned} \vec{e}_2 &= \left(1 + \frac{\langle \vec{e}_1, \vec{e}_3 \rangle}{\|\vec{e}_1\|^2}\right) \vec{e}_2 \\ &= \begin{bmatrix} \tilde{e}_2 \\ 0 \end{bmatrix}. \end{aligned}$$

To preserve  $JH_\omega \vec{e}_2 = \vec{e}_1$ , we get that  $\vec{e}_1$  must be replaced by

$$\begin{aligned} \vec{e}_1 &= \left(1 + \frac{\langle \vec{e}_1, \vec{e}_3 \rangle}{\|\vec{e}_1\|^2}\right) \vec{e}_1 \\ &= \begin{bmatrix} 0 \\ \tilde{e}_1 \end{bmatrix}. \end{aligned}$$

With these substitutions, the  $JH_\omega$  matrix on  $D_\omega$  remains the same and we obtain the relationship  $\langle \vec{e}_3, \vec{e}_1 \rangle = 0$ . From here on we will assume that we are working with this modified basis and simply take  $\vec{e}_j := \tilde{e}_j$  for  $j = 1, 2, 3, 4$ .

We will define  $\vec{\xi}_i$  to be the dual basis to the revised  $\vec{e}_i$  within  $D_\omega$ . That is, the  $\vec{\xi}_i$  are defined by  $\vec{\xi}_i \in D_\omega$  and

$$\langle \vec{\xi}_i, \vec{e}_j \rangle = \delta_{ij}.$$

If we make the change of basis described above, then we can compute the  $\vec{\xi}_j$  as follows. Define  $D = \|\vec{e}_2\|^2\|\vec{e}_4\|^2 - \langle \vec{e}_2, \vec{e}_4 \rangle^2$ . Then:

$$\begin{aligned}\vec{\xi}_1 &= \frac{1}{\|\vec{e}_1\|^2} \vec{e}_1, \\ \vec{\xi}_2 &= \frac{\|\vec{e}_4\|^2}{D} \vec{e}_2 - \frac{\langle \vec{e}_2, \vec{e}_4 \rangle}{D} \vec{e}_4, \\ \vec{\xi}_3 &= \frac{1}{\|\vec{e}_3\|^2} \vec{e}_3, \\ \vec{\xi}_4 &= -\frac{\langle \vec{e}_2, \vec{e}_4 \rangle}{D} \vec{e}_2 + \frac{\|\vec{e}_2\|^2}{D} \vec{e}_4.\end{aligned}$$

As with the  $\vec{e}_j$ 's,

$$\vec{\xi}_j = \begin{bmatrix} 0 \\ \xi_j \end{bmatrix}$$

for  $j = 1, 3$  and

$$\vec{\xi}_j = \begin{bmatrix} \xi_j \\ 0 \end{bmatrix}$$

for  $j = 2, 4$  to distinguish between vectors and their scalar components.

### 3. DERIVATION OF THE ODES

To derive the ODEs we start with a small spherically symmetric perturbation of the minimal mass soliton, then project onto the discrete spectral subspace. Here, we closely follow [20].

We begin with the following ansatz, which allows *theta* and  $\omega$  to modulate:

$$(3.1) \quad \vec{u}(t) = e^{(J_0^t \omega(t') dt' + \theta(t))J} (\vec{\phi}_{\omega(t)} + \vec{\rho}(t)).$$

Here we are suppressing the dependence of  $u$ ,  $\phi$ , and  $\rho$  on  $|x|$  for notational simplicity. Recall, we have assumed  $u$  to be spherically symmetric, so no other modulation parameters occur. Unlike in [20] we do not assume that the rotation variable  $\theta(t)$  is identically zero, so we need to include  $\theta$  modulation in our full ansatz. Note that the derivation which follows applies for all nonlinearities in any dimension; specialization is required only to get explicit numerical results. This model includes the full dynamical system for spherically symmetric data and is designed in such a way that coupling to the continuous spectral subspace could easily be reintroduced. The authors plan to analyze the effect of that coupling, which is expected to be dissipative, in a future work.

Differentiating (3.1) with respect to  $t$ , we get

$$\vec{u}_t = [(\omega + \dot{\theta})J(\vec{\phi} + \vec{\rho}) + \dot{\omega}\vec{\phi}' + \dot{\vec{\rho}}] e^{i(J_0^t \omega(t') dt' + \theta(t))},$$

where for a time dependent function  $g(t)$  we represent differentiation with respect to  $t$  by  $\partial_t g = \dot{g}$  and for an  $\omega$  dependent function  $f(\omega)$  we denote differentiation with respect to the soliton parameter  $\omega$  by  $\partial_\omega f = f'$ . Plugging the above ansatz into equation (1.1) and cancelling the phase term yields

$$(3.2) \quad -(\omega + \dot{\theta})(\vec{\phi} + \rho) + \dot{\omega}J\vec{\phi}' + J\dot{\rho} + \Delta\vec{\phi} + \Delta\vec{\rho} + g(|\phi + \rho|^2)(\vec{\phi} + \vec{\rho}) = 0.$$

Recall that since  $\phi$  is a soliton solution,  $-\omega\phi + \Delta\phi + g(|\phi|^2)\phi = 0$ , yielding

$$(3.3) \quad -\dot{\theta}\vec{\phi} - (\omega + \dot{\theta})(\vec{\rho}) + \dot{\omega}J\vec{\phi}' + J\dot{\rho} + \Delta\vec{\rho} + g(|\vec{\phi} + \vec{\rho}|^2)(\vec{\phi} + \vec{\rho}) - g(\phi^2)\phi = 0.$$

We multiply by  $J$ , solve for  $\vec{\rho}$ , and simplify. At the same time, we collect the  $\Delta\vec{\rho}$  and  $-\omega\vec{\rho}$  terms with the linear portion of  $g(|\vec{\phi} + \vec{\rho}|^2)(\vec{\phi} + \vec{\rho}) - g(\phi^2)\vec{\phi}$ , which yields  $JH_\omega$  as defined in (2.5). The remaining terms of the nonlinearity are at least quadratic in  $\rho$ ; recall that the quadratic terms are described in (2.3) and denoted  $N(\omega, \rho)$ .

Defining  $\rho_j(t)$  as the coefficient of  $\vec{e}_j(t)$  in  $\rho$ , we have

$$\vec{\rho} = \begin{bmatrix} \rho_{Re} \\ \rho_{Im} \end{bmatrix} = \rho_1\vec{e}_1 + \rho_2\vec{e}_2 + \rho_3\vec{e}_3 + \rho_4\vec{e}_4 + \vec{\rho}_c.$$

Then, the above calculations give us

$$(3.4) \quad \dot{\vec{\rho}} = JH_\omega\vec{\rho} - \dot{\theta} \begin{pmatrix} 0 \\ \phi \end{pmatrix} - \dot{\theta}J\vec{\rho} - \dot{\omega} \begin{pmatrix} \phi' \\ 0 \end{pmatrix} + \vec{N}(\omega, \vec{\rho}).$$

Taking the inner product of (3.4) with each of the  $\vec{\xi}_i$  as defined in Section 2.2, and applying (2.8) yields the following system:

$$(3.5) \quad \begin{aligned} \langle \vec{\xi}_1, \dot{\vec{\rho}} \rangle &= \rho_2 - \dot{\theta} - \dot{\theta}\langle \vec{\xi}_1, J\vec{\rho} \rangle + \langle \vec{\xi}_1, \vec{N} \rangle, \\ \langle \vec{\xi}_2, \dot{\vec{\rho}} \rangle &= \rho_3 - \dot{\omega} - \dot{\theta}\langle \vec{\xi}_2, J\vec{\rho} \rangle + \langle \vec{\xi}_2, \vec{N} \rangle, \\ \langle \vec{\xi}_3, \dot{\vec{\rho}} \rangle &= \rho_4 - \dot{\theta}\langle \vec{\xi}_3, J\vec{\rho} \rangle + \langle \vec{\xi}_3, \vec{N} \rangle, \\ \langle \vec{\xi}_4, \dot{\vec{\rho}} \rangle &= a(\omega)\rho_3 - \dot{\theta}\langle \vec{\xi}_4, J\vec{\rho} \rangle + \langle \vec{\xi}_4, \vec{N} \rangle. \end{aligned}$$

From this point forward in our approximation we drop the  $\vec{\rho}_c$  component as a higher order error term. Using the product rule, we solve the left hand side for  $\dot{\rho}_i$  and put the extra terms from the derivative of the operator that projects onto the discrete spectral subspace onto the right hand side.

We have, as in [20], that

$$P_d\dot{\vec{\rho}} = \sum \vec{e}_j\dot{\rho}_j + \dot{\omega} \sum \vec{e}_i\Gamma_{ij}\rho_j - \dot{\omega}P_dP_d'\vec{\rho},$$

where we have implicitly defined

$$\Gamma_{ij} = \langle \xi_i, e_j' \rangle$$

and used that

$$P_d \frac{d}{dt} P_c \vec{\rho} = -\dot{\omega} P_d P_d' \vec{\rho}.$$

This gives

$$(3.6) \quad \begin{aligned} \dot{\rho}_1 + \dot{\theta} &= \rho_2 - \dot{\theta} \langle \vec{\xi}_1, J\vec{\rho} \rangle + \langle \vec{\xi}_1, \vec{N} \rangle + \dot{\omega} (\langle \vec{\xi}_1, P_d' \vec{\rho} \rangle - \sum \Gamma_{1j} \rho_j), \\ \dot{\rho}_2 + \dot{\omega} &= \rho_3 - \dot{\theta} \langle \vec{\xi}_2, J\vec{\rho} \rangle + \langle \vec{\xi}_2, \vec{N} \rangle + \dot{\omega} (\langle \vec{\xi}_2, P_d' \vec{\rho} \rangle - \sum \Gamma_{2j} \rho_j), \\ \dot{\rho}_3 &= \rho_4 - \dot{\theta} \langle \vec{\xi}_3, J\vec{\rho} \rangle + \langle \vec{\xi}_3, \vec{N} \rangle + \dot{\omega} (\langle \vec{\xi}_3, P_d' \vec{\rho} \rangle - \sum \Gamma_{3j} \rho_j), \\ \dot{\rho}_4 &= a(\omega) \rho_3 - \dot{\theta} \langle \vec{\xi}_4, J\vec{\rho} \rangle + \langle \vec{\xi}_4, \vec{N} \rangle + \dot{\omega} (\langle \vec{\xi}_4, P_d' \vec{\rho} \rangle - \sum \Gamma_{4j} \rho_j). \end{aligned}$$

There is also coupling to the continuous spectrum through terms such as  $\langle \vec{\xi}_1, J\vec{\rho} \rangle$  which we omit. This can be included in the error term and is not analyzed in our finite dimensional system.

To make the system well-determined, we must introduce two orthogonality conditions. The first is  $\langle \rho, e_2 \rangle = 0$ , and the second is  $\langle \rho, e_1 \rangle = 0$ . These represent the choice of  $\omega(t)$  and  $\theta(t)$  respectively that minimize the size of  $\rho$ , which yields  $\rho_2 = \dot{\rho}_2 = 0$ , and  $\rho_1 = \dot{\rho}_1 = 0$ , respectively.

The reduced system is then:

$$(3.7) \quad \begin{aligned} \dot{\theta} &= -\dot{\theta} \langle \vec{\xi}_1, J\vec{\rho} \rangle + \langle \vec{\xi}_1, \vec{N} \rangle + \dot{\omega} (\langle \vec{\xi}_1, P_d' \vec{\rho} \rangle - \sum \Gamma_{1j} \rho_j), \\ \dot{\omega} &= \rho_3 - \dot{\theta} \langle \vec{\xi}_2, J\vec{\rho} \rangle + \langle \vec{\xi}_2, \vec{N} \rangle + \dot{\omega} (\langle \vec{\xi}_2, P_d' \vec{\rho} \rangle - \sum \Gamma_{2j} \rho_j), \\ \dot{\rho}_3 &= \rho_4 - \dot{\theta} \langle \vec{\xi}_3, J\vec{\rho} \rangle + \langle \vec{\xi}_3, \vec{N} \rangle + \dot{\omega} (\langle \vec{\xi}_3, P_d' \vec{\rho} \rangle - \sum \Gamma_{3j} \rho_j), \\ \dot{\rho}_4 &= a(\omega) \rho_3 - \dot{\theta} \langle \vec{\xi}_4, J\vec{\rho} \rangle + \langle \vec{\xi}_4, \vec{N} \rangle + \dot{\omega} (\langle \vec{\xi}_4, P_d' \vec{\rho} \rangle - \sum \Gamma_{4j} \rho_j). \end{aligned}$$

In [20], the authors further reduce this system to prove there is an initial nonlinear instability. (Note that they have a slightly different system because they have assumed that  $\theta \equiv 0$ .) We are interested in the dynamics on an intermediate time scale; thus, we retain quadratically nonlinear terms in our equations.

Our notation is as follows. First, we have

$$\begin{aligned} \langle \vec{\xi}_1, J\vec{\rho} \rangle &= \langle \xi_1, \rho_2 \phi' + \rho_4 \beta \rangle \\ &= \rho_4 \langle \xi_1, \beta \rangle, \end{aligned}$$

since  $\rho_2 = 0$ . Denote

$$c_{14} = \langle \xi_1(\omega^*), \beta(\omega^*) \rangle,$$

which is the highest order term and the only one that will figure into our quadratic expansion. Notice that this is a real inner product of functions that normally do not appear in the same component of the complex vectors, because of the  $J$  in the equation.

Similarly, we have

$$\begin{aligned}\langle \vec{\xi}_2, J\vec{\rho} \rangle &= \langle \xi_2, -\rho_1\phi - \rho_3\alpha \rangle \\ &= -\rho_3\langle \xi_2, \alpha \rangle,\end{aligned}$$

since  $\rho_1 = 0$ . Denote

$$c_{23} = \langle \xi_2(\omega^*), \alpha(\omega^*) \rangle,$$

which is again the highest order term.

Then we have

$$\begin{aligned}\langle \vec{\xi}_3, J\vec{\rho} \rangle &= \langle \xi_3, \rho_2\phi' + \rho_4\beta \rangle \\ &= \rho_4\langle \xi_3, \beta \rangle,\end{aligned}$$

since  $\rho_2 = 0$ . Denote

$$c_{34} = \langle \xi_3(\omega^*), \beta(\omega^*) \rangle.$$

Finally, we have

$$\begin{aligned}\langle \vec{\xi}_4, J\vec{\rho} \rangle &= \langle \xi_4, -\rho_1\phi - \rho_3\alpha \rangle \\ &= -\rho_3\langle \xi_4, \alpha \rangle,\end{aligned}$$

since  $\rho_1 = 0$ . Denote by

$$c_{43} = \langle \xi_4(\omega^*), \alpha(\omega^*) \rangle.$$

We also write  $g_{ij}$  for the term  $\Gamma_{ij}(\omega) = \langle \vec{\xi}_i, \vec{e}_j \rangle$  at  $\omega = \omega^*$ .

Next, consider the terms  $\langle \vec{\xi}_j, P'_d\vec{\rho} \rangle$ . These terms are the  $e_j$  components of  $P'_d\vec{\rho}$ . We have:

$$\begin{aligned}P_d P'_d \vec{\rho} &= P_d \left[ \sum_{j=1}^4 \langle \vec{\xi}_j, \vec{\rho} \rangle \vec{e}_j + \sum_{j=1}^4 \langle \vec{\xi}_j, \vec{\rho} \rangle \vec{e}'_j \right] \\ &= \sum_{j=1}^4 \sum_{k=3}^4 \langle \vec{\xi}_j, \vec{e}_k \rangle \rho_k \vec{e}_j + \rho_3 P_d \vec{e}'_3 + \rho_4 P_d \vec{e}'_4 \\ &= \sum_{j=1}^4 \sum_{k=3}^4 \langle \vec{\xi}_j, \vec{e}_k \rangle \rho_k \vec{e}_j + \rho_3 (\Gamma_{13} \vec{e}_1 + \Gamma_{33} \vec{e}_3) + \rho_4 (\Gamma_{24} \vec{e}_2 + \Gamma_{44} \vec{e}_4) \\ &= \langle \vec{\xi}_1, \vec{e}_3 \rangle \rho_3 \vec{e}_1 + \langle \vec{\xi}_2, \vec{e}_4 \rangle \rho_4 \vec{e}_2 + \langle \vec{\xi}_3, \vec{e}_3 \rangle \rho_3 \vec{e}_3 + \langle \vec{\xi}_4, \vec{e}_4 \rangle \rho_4 \vec{e}_4 \\ &\quad + \rho_3 (\Gamma_{13} \vec{e}_1 + \Gamma_{33} \vec{e}_3) + \rho_4 (\Gamma_{24} \vec{e}_2 + \Gamma_{44} \vec{e}_4) \\ &= (\langle \vec{\xi}_1, \vec{e}_3 \rangle + \Gamma_{13}) \rho_3 \vec{e}_1 + (\langle \vec{\xi}_2, \vec{e}_4 \rangle + \Gamma_{24}) \rho_4 \vec{e}_2 \\ &\quad + (\langle \vec{\xi}_3, \vec{e}_3 \rangle + \Gamma_{33}) \rho_3 \vec{e}_3 + (\langle \vec{\xi}_4, \vec{e}_4 \rangle + \Gamma_{44}) \rho_4 \vec{e}_4.\end{aligned}$$

Therefore, the relevant nonzero terms are

$$\begin{aligned}\langle \vec{\xi}_1, P'_d \vec{\rho} \rangle &= (\langle \vec{\xi}_1^T, \vec{e}_3 \rangle + \Gamma_{13}) \rho_3, \\ \langle \vec{\xi}_2, P'_d \vec{\rho} \rangle &= (\langle \vec{\xi}_2^T, \vec{e}_4 \rangle + \Gamma_{24}) \rho_4.\end{aligned}$$

We denote

$$\begin{aligned}p_{13} &= \langle \vec{\xi}_1^T(\omega^*), \vec{e}_3(\omega^*) \rangle, \\ p_{33} &= \langle \vec{\xi}_3^T(\omega^*), \vec{e}_3(\omega^*) \rangle,\end{aligned}$$

and

$$\begin{aligned}p_{24} &= \langle \vec{\xi}_2^T(\omega^*), \vec{e}_4(\omega^*) \rangle, \\ p_{44} &= \langle \vec{\xi}_4^T(\omega^*), \vec{e}_4(\omega^*) \rangle.\end{aligned}$$

Note that some cancellation will occur with the  $\Gamma_{ij}$  terms that appear separately in the system of ODEs, leaving only these  $p_{ij}$  terms in the finally system.

Finally, the terms  $\langle \vec{\xi}_i, \vec{N}(\omega, \vec{\rho}) \rangle$  must be computed. We are only interested in the quadratic terms, which, according to (2.3) are:

$$(3.8) \quad 3Jg'(\phi^2)\phi\rho_{Re}^2 + 2Jg''(\phi^2)\phi^2\rho_{Re}^2 + Jg'(\phi^2)\phi\rho_{Im}^2 + 2g'(\phi^2)\phi\rho_{Re}\rho_{Im}.$$

Recall that, since  $\rho_1$  and  $\rho_2$  are 0, the projection onto the discrete-spectrum of  $\rho_{Re}$  is just  $\rho_3\vec{e}_3$  and the projection onto the discrete-spectrum of  $\rho_{Im}$  is just  $\rho_4\vec{e}_4$ . We now have to compute the lowest-order terms of

$$\langle \vec{\xi}_1, \vec{N}(\omega, \vec{\rho}) \rangle.$$

The multiplier of  $\rho_3^2$  in  $\langle \vec{\xi}_1, \vec{N}(\omega, \vec{\rho}) \rangle$  is

$$n_{133} = \langle \xi_1, (3g'(\phi^2)\phi + 2g''(\phi^2)\phi^2)e_3^2 \rangle.$$

Similarly, we define

$$\begin{aligned}n_{144} &= \langle \xi_1, g'(\phi^2)\phi e_4^2 \rangle, \\ n_{234} &= \langle \xi_2, -2g'(\phi^2)\phi e_3 e_4 \rangle, \\ n_{333} &= \langle \xi_3, (3g'(\phi^2)\phi + 2g''(\phi^2)\phi^2)e_3^2 \rangle, \\ n_{344} &= \langle \xi_3, g'(\phi^2)\phi e_4^2 \rangle, \\ n_{434} &= \langle \xi_4, -2g'(\phi^2)\phi e_3 e_4 \rangle.\end{aligned}$$

Notice that, as in the computation of the  $c_{ij}$ , these are real inner products between real functions that normally appear in different components of the complex vectors.

Lastly, we need to estimate  $a(\omega)$ . Recall that  $a(\omega^*) = 0$ , and that  $a(\omega)$  appears in (3.7) multiplied by  $\rho_3$ , so we are seeking only the linear term,  $a(\omega) \sim a_0(\omega - \omega^*)$ . We calculate:

$$a_0 = a'(\omega^*) = -\frac{2}{\langle \phi_{\omega^*}, \beta \rangle} (\langle \phi'_{\omega^*}, \phi'_{\omega^*} \rangle - \langle \phi_{\omega^*}, \phi''_{\omega^*} \rangle).$$

With these assumptions, we conclude the following:

**Proposition 3.1.** *The quadratic approximation for the evolution of a perturbation of the minimal mass soliton, (3.3), ignoring coupling to the continuous spectrum, is*

$$\begin{aligned} \dot{\theta} &= -c_{14}\dot{\theta}\rho_4 + n_{133}\rho_3^2 + n_{144}\rho_4^2 + \dot{\omega}p_{13}\rho_3, \\ \dot{\omega} &= \rho_3 + c_{23}\dot{\theta}\rho_3 + n_{234}\rho_3\rho_4 + \dot{\omega}p_{24}\rho_4, \\ (3.9) \quad \dot{\rho}_3 &= \rho_4 - c_{34}\dot{\theta}\rho_4 + n_{333}\rho_3^2 + n_{344}\rho_4^2 + \dot{\omega}p_{33}\rho_3, \\ \dot{\rho}_4 &= a_0(\omega - \omega^*)\rho_3 + c_{43}\dot{\theta}\rho_3 + n_{434}\rho_3\rho_4 + \dot{\omega}p_{44}\rho_4. \end{aligned}$$

In this system we implicitly assume that  $\theta$ ,  $(\omega - \omega^*)$ ,  $\rho_i$  and their time derivatives are all of the same order.

#### 4. NUMERICAL METHODS

From here on, we use numerical techniques to analyze solutions to (3.9). We will work in one space dimension, with the specific saturated nonlinearity  $g(s) = \frac{s^3}{1+s^2}$  as described in the introduction. In this setting, our assumption of spherical symmetry on the initial data becomes an assumption that  $u_0$  is even. Though we have a complete description of the generalized kernel of  $J\mathcal{H}$ , including its size and the relation among the elements, nothing is expressible in terms of elementary functions. As this kernel determines the coefficients in our ODE system, we numerically compute it, permitting us to subsequently integrate the ODEs numerically.

The sinc function,  $\sin(\pi x)/(\pi x)$  was used to compute solitary wave solutions when analytical expressions were not readily available in [39] and in the forthcoming [52]. It has also been used to study time dependent nonlinear wave equations, [5, 46, 11, 18], and a variety of linear and nonlinear boundary value problems, [13, 14, 26, 27, 25, 42].

We will use the sinc function to estimate the coefficients in three steps:

- Compute a discrete representation of the minimal mass soliton,  $\phi_{\omega^*}$ .
- Compute discrete representations of the generalized kernel of  $\mathcal{H}$ , i.e. the derivatives with respect to  $\omega$ .
- Compute necessary inner products for the coefficients.



**4.1. Sinc Discretization.** The problem of finding a soliton solution of (1.5) is a nonlinear boundary value problem posed on  $\mathbb{R}$ . We respect this description in our discretization by approximating functions with the sinc spectral method. This technique is thoroughly explained in [38, 56, 55, 57] and briefly in Appendix A.1. In the sinc discretization, the problem remains posed on  $\mathbb{R}$  and the boundary conditions, that the solution vanish at  $\pm\infty$ , are naturally incorporated.

Given a function  $u(x) : \mathbb{R} \rightarrow \mathbb{R}$ ,  $u$  is approximated using a superposition of shifted and scaled sinc functions:

$$(4.1) \quad C_{M,N}(u, h)(x) \equiv \sum_{k=-M}^N u_k \operatorname{sinc}\left(\frac{x - x_k}{h}\right) = \sum_{k=-M}^N u_k S(k, h)(x),$$

where  $x_k = kh$  for  $k = -M, \dots, N$  are the *nodes* and  $h > 0$ . There are *three* parameters in this discretization,  $h$ ,  $M$ , and  $N$ , determining the number of and spacing of lattice points. This is common to numerical methods posed on unbounded domains; see [15].

A useful and important feature of this spectral method is that, when evaluated at a node,

$$(4.2) \quad C_{M,N}(u, h)(x_k) = u_k.$$

Additionally, the convergence is rapid both in practice and theoretically. See Theorem 1 in Appendix A.1 for a statement on optimal convergence.

Since the soliton is an even function, we may take  $N = M$ . We will thus write

$$(4.3) \quad C_M(u, h)(x) \equiv C_{M,M}(u, h)(x).$$

The symmetry implies  $u_{-k} = u_k$  for  $k = -M, \dots, M$ . We take advantage of this constraint in our computations. In addition, we slave  $h$  to  $M$  in accordance with (A.8).

To compute a discrete sinc approximation of the ground state, we frame the soliton equation as a *nonlinear collocation problem*. Approximating  $\phi(x)$  as in (4.1), we seek coefficients  $\{R_k\}$  such that

$$(4.4) \quad \begin{aligned} & \partial_x^2 C_M(\phi, h)(x_k) - \lambda C_M(\phi, h)(x_k) \\ & + g(|C_M(\phi, h)(x_k)|^2) C_M(\phi, h)(x_k) = 0, \quad \text{for } k = -M, \dots, M. \end{aligned}$$

By satisfying (4.4), the discrete approximation solves the soliton equation in the *strong* sense at the nodes, also known as *collocation points*. This is in contrast to a Galerkin formulation, which solves the equation in the weak sense. However, for the one dimensional under consideration, sinc-Galerkin and sinc-collocation lead to the same algebraic system.

(4.4) yields a system of nonlinear algebraic equations. Let  $\vec{\phi}$  be the column vector associated with the discrete approximation of  $\phi$ :

$$(4.5) \quad C_M(\phi, h)(x_k) \mapsto \vec{\phi} = \begin{pmatrix} \phi_{-M} \\ \phi_{-M+1} \\ \vdots \\ \phi_M \end{pmatrix}.$$

Differentiation of a sinc approximated function that is evaluated at the collocation points corresponds to matrix multiplication:

$$(4.6) \quad \partial_x^2 C_M(\phi, h)(x_k) \mapsto D^{(2)} \vec{\phi}.$$

Explicitly,  $D^{(2)}$  is

$$(4.7) \quad D_{jk}^{(2)} = \frac{d^2}{dx^2} S(j, h)(x)|_{x=x_k} = \begin{cases} \frac{1}{h^2} \frac{-\pi^2}{3} & j = k \\ \frac{1}{h^2} \frac{-2(-1)^{k-j}}{(k-j)^2} & j \neq k \end{cases}.$$

Using 4.2,

$$g(|C_M(\phi, h)(x_k)|^2) C_M(\phi, h)(x_k) = C_M(g(|\phi|^2)\phi, h)(x_k) = g(|\phi_k|^2)\phi_k.$$

Thus

$$g(|C_M(\phi, h)(x_k)|^2) C_M(\phi, h)(x_k) \mapsto g(|\vec{\phi}|^2) \vec{\phi} \equiv \begin{pmatrix} g(|\phi_{-M}|^2)\phi_{-M} \\ \vdots \\ g(|\phi_M|^2)\phi_M \end{pmatrix}.$$

With these relations, the discrete system is

$$(4.8) \quad D^{(2)} \vec{\phi} - \omega \vec{\phi} + g(|\vec{\phi}|^2) \vec{\phi} = 0.$$

It is this equation to which we apply a nonlinear solver, subject to an appropriate guess. We discuss an important subtlety in Appendix A.2.

**4.2. Computing the Minimal Mass.** Now that we have an algorithm for finding a discrete representation of a soliton, we seek to find the value of the soliton parameter for the one possessing minimal mass, along with the corresponding discretized soliton. The sinc discretization has the property that the  $L^2(\mathbb{R})$  inner product is well approximated by

$$\langle f, g \rangle = \int f g dx \approx \sum_{-M}^M h f_k g_k = h(\vec{f} \cdot \vec{g}).$$

Thus, the mass of the soliton can be estimated by

$$\|\phi\|_{L^2}^2 \approx h|\vec{\phi}|^2.$$

Recognizing that  $\vec{\phi} = \vec{\phi}(\omega)$ , we seek to minimize the functional  $h|\vec{\phi}(\omega)|^2$  with respect to  $\omega$ . The argument  $\omega$  for which the minimum occurs will be  $\omega^*$ . To find the minimal mass,

we take the derivative, getting a discrete representation of the minimal mass orthogonality condition:

$$(4.9) \quad 2h\vec{\phi} \cdot \vec{\phi}' = 0.$$

We solve (4.9) to find  $\omega^*$ , computing  $\vec{\phi}_{\omega^*}$  in the process.

The value  $\omega^*$  can be obtained by other algorithms. In the one-dimensional case, the soliton equation possesses a first integral, permitting the minimal mass to be computed by numerical quadrature and minimization; a comparison of our results and this approach appears in Appendix A.4.1. Though these approaches are quite accurate for the task of computing the minimal mass, they are inadequate at computing the generalized kernel. Thus, we seek to solve the problem consistently by finding the minimal mass for a given  $2M + 1$  dimensional approximation of the problem.

**4.3. Discretized Generalized Kernel.** As seen in Section 2.2, at the minimal mass soliton  $\phi_{\omega^*}$ , there are four functions associated with the kernel satisfying the second order equations:

$$(4.10) \quad L_- \phi_{\omega^*} = 0,$$

$$(4.11) \quad L_+(-\phi'_{\omega^*}) = \phi_{\omega^*},$$

$$(4.12) \quad L_- \alpha = -\phi'_{\omega^*},$$

$$(4.13) \quad L_+ \beta = \alpha.$$

These four functions can also be discretized with sinc, as in (4.1). The operators,  $L_{\pm}$ , have discrete spectral representations:

$$(4.14) \quad L_+ \mapsto \mathbf{L}_+ \equiv -D^{(2)} + \omega I - \text{diag}\{g(\vec{\phi}_{\omega^*}^2)\},$$

$$(4.15) \quad L_- \mapsto \mathbf{L}_- \equiv -D^{(2)} + \omega I - \text{diag}\{g(\vec{\phi}_{\omega^*}^2) - 2g'(\vec{\phi}_{\omega^*}^2)\vec{\phi}_{\omega^*}^2\}.$$

Taking  $\vec{u} = \vec{\phi}$ , we successively solve for  $\vec{\phi}'$ ,  $\vec{\alpha}$ , and  $\vec{\beta}$ . A singular value decomposition must be used to get  $\vec{\alpha}$  since  $\mathbf{L}_-$  has a non-trivial kernel.

Furthermore, we compute discrete approximations of the derivatives of  $\phi$ ,  $\phi'$ ,  $\alpha$ , and  $\beta$  taken with respect to  $\omega$  at  $\omega^*$ . The relevant operators are formed analogously to (4.14) and (4.15).

**4.4. Convergence.** Amongst the many calculations made, the most important is of  $\omega^*$ , the parameter of the minimal mass soliton. We summarize the results in Table 1. We see that  $h|\vec{\phi}|^2$  robustly converges, achieving twelve digits of precision and  $\omega^*$  appears to achieve eleven digits of precision. These are consistent with the values in Table 4 from Appendix A.4, where they were computed using a different methods.

For the purposes of our simulations, we believe we have sufficient precision, approximately ten significant digits, for the time integration of our system of ODEs. Some data for the convergence of the coefficients appearing in (3.9) is given in Appendix A.3.

TABLE 1. The convergence of the sinc discretization to the minimal mass soliton.

$M$	$h \vec{\phi} ^2$	$\omega^*$
20	3.820771417633398	0.177000229690401
40	3.821145471868853	0.177576993694258
60	3.821148930202135	0.177587655985074
80	3.821149018422933	0.177588043323139
100	3.821149022493814	0.177588063805561
200	3.821149022780618	0.177588065432740
300	3.821149022780439	0.177588065432795
400	3.821149022780896	0.177588065433095
500	3.821149022780275	0.177588065432928

## 5. NUMERICAL RESULTS

We explore here the dynamics of the finite dimensional system (3.9) and compare with solutions for the full nonlinear PDE (1.1) with corresponding initial data.

To solve (3.9), we use the solver *ode45* from MATLAB after properly preparing the initial data using the soliton finding codes in Section 4.1.

**5.1. PDE Solver.** In order to determine the accuracy of our results, we also use a nonlinear solver to approximate the solutions with a perturbed minimal-mass soliton as initial data. For this nonlinear solver, we use a finite element scheme in space and a Crank-Nicholson scheme in time. This is similar to the method used in [30]. In brief, we discretize our (1.1) by method of lines, using finite elements in space and Crank-Nicholson for time-stepping. This method is  $L^2$  conservative, though it is not energy conserving. A similar scheme was implemented without potential in [4], where the blow-up for NLS in several dimensions was investigated.

We require the spatial grid to be large enough to ensure negligible interaction with the boundary. As absorbing boundary conditions for cubic NLS currently require high frequency limits to apply successfully, we choose simply to carefully ensure that our grid is large enough in order for the interactions to be negligible throughout the experiment. For the convergence of such methods without potentials see the references in [4], [2] and [3].

We select a symmetric region about the origin,  $[-R, R]$ , upon which we place a mesh of  $N$  elements. The standard hat function basis is used in the Galerkin approximation. We allow for a finer grid in a neighbourhood of length 1 centered at the origin to better study the effects of the soliton interactions.

In terms of the hat basis the PDE (1.1) becomes:

$$\begin{aligned} \langle u_t, v \rangle + i \langle u_x, v_x \rangle / 2 - i \langle g(|u|^2)u, v \rangle &= 0, \\ u(0, x) = u_0, \quad u(t, x) &= \sum_v c_v(t) v, \end{aligned}$$

where  $\langle \cdot, \cdot \rangle$  is the standard  $L^2$  inner product,  $v$  is a basis function and  $u, u_0$  are linear combinations of the  $v$ 's.

Since the  $v$ 's are hat functions, we have a tridiagonal linear system. Let  $h_t > 0$  be a uniform time step, and let

$$u_n = \sum_v c_v(nh_t) v$$

be the approximate solution at the  $n$ -th time step. Implementing Crank-Nicholson, the system becomes:

$$\begin{aligned} \langle u_{n+1} - u_n, v \rangle + ih_t \langle ((u_{n+1} + u_n)/2)_x, v_x \rangle \\ = ih_t \langle g(|(u_{n+1} + u_n)/2|^2)(u_{n+1} + u_n)/2, v \rangle, \quad u_0 = \sum_v \alpha_v v. \end{aligned}$$

By defining

$$y_n = (u_{n+1} + u_n)/2,$$

we have simplified our system to:

$$\langle y_n, v \rangle + i \frac{h_t}{4} \langle (y_n)_x, v_x \rangle = i \frac{h_t}{2} \langle |y_n|^2 y_n, v \rangle + \langle u_n, v \rangle.$$

An iteration method from [4] is now used to solve this nonlinear system of equations. Namely, we set,

$$\langle y_n^{k+1}, v \rangle + i \frac{h_t}{4} \langle (y_n^{k+1})_x, v_x \rangle = i \frac{h_t}{2} \langle |y_n^k|^2 y_n^k, v \rangle + \langle u_n, v \rangle.$$

We take  $y_n^0 = u_n$  and perform three iterations in order to obtain an approximate solution.

For our problem, we have taken (1.1) with the nonlinearity

$$\frac{|u|^6}{1 + |u|^4} u.$$

Then, the minimal mass soliton occurs at

$$\omega^* = .177588065433.$$




FIGURE 2. A plot of the maximum amplitude with respect to the  $L^2$  norm for a saturated nonlinear Schrödinger equation. Computed at  $M + 1 = 101$  collocation points for  $\omega \in [0.01, 1.5]$ .

**5.2. Results.** With the numerical schemes outlined above, we then compare our finite dimensional model to the numerically integrated solution with appropriate initial data  $\omega_0 = \omega(0) < \omega^*$ ,  $\alpha_0 = \rho_3(0)$ ,  $\beta_0 = \rho_4(0)$  and  $\theta_0 = \theta(0) = 0$  for simplicity. In Figures 3, 4, 5, we take  $\beta_0 > 0$  and vary  $\alpha_0, \omega_0$ . Similarly, in Figures 6, 7, 8, we take  $\beta_0 < 0$  and once again vary  $\alpha_0, \omega_0$ . Note that we are comparing solutions to the ODEs to solutions of (1.1) with the correct initial parameters so that the initial profiles are identical. The initial data for both the finite and infinite dimensional solvers always consists of an orbitally unstable soliton being perturbed by elements of the generalized kernel of the linearized operator  $\mathcal{H}$  at  $\omega^*$ . Also, all the numerics we present are given in terms of plotting  $|u(t, 0)|$ , the amplitude of the solution at 0, versus time  $t$ . The computed correlation between the amplitude of a soliton  $\phi_\omega$  at 0 and the soliton parameter  $\omega$  can be seen in Figure 2.

When  $\beta_0 > 0$ , the initial data is expected to allow the admission of a solution with a soliton component as  $t$  increases. The finite dimensional system shows that if we initially perturb the system either towards the stable or the unstable side of the curve, the system produces immediate oscillations. Specifically, if the dynamics begin to diverge, the higher order nonlinear corrections in (3.9) arrest the solution, resulting in fairly uniform oscillations about the minimal mass soliton. See Figures 3, 4, 5. As one can see, for initial values  $\omega_0 \approx \omega^*$ , we see a good fit for several oscillations of our finite dimensional approximation to the dynamics of the full solution. As expected, this weakens as  $\omega_0$  diverges from  $\omega^*$  due to the nature of our approximations in Section 3. We conjecture that such oscillations

about the minimal mass when coupled to the continuous spectrum will lead generically to a damped convergence of the solution towards the minimal mass soliton on a long time scale, similar to the oscillations observed in [37].

For  $\beta_0 < 0$ , and other initial parameters sufficiently small, the initial data is below the minimal mass and is expected to disperse as  $t \rightarrow \infty$ . In this regime, we see the amplitude fall below the value associated with the minimal mass soliton, forcing  $\omega(t) < \omega_*$ , the minimal mass value. Thus, we are on the unstable branch of the soliton curve. Clearly, conservation of mass for (1.1) forbids the primitive system from indefinitely behaving as a perturbed soliton with a progressively smaller value of  $\omega$ . Indeed, we see divergence of the nonlinear solution from our solution as  $t$  increases. As our approximation is based on an expansion about  $\omega_*$ , the deviation eventually renders it invalid.

However, on shorter time scales there is reasonable agreement between the full solution and our finite dimensional approximation; see Figures 6, 7, 8. This occurs regardless of whether we initially perturb in the stable or unstable direction. Our findings confirm the regimes predicted in [43].

It remains to briefly comment on the convergence of our numerical methods. The ODE solver, *ode45*, for the finite dimensional system of ODEs is a standard Runge-Kutta method with known strong convergence results documented in a number of introductory texts on numerical methods. In addition, the finite element solver for the full nonlinear problem has well-established analytic convergence results, see [2]. Hence, the solutions for the corresponding systems are known to be accurate representations of the actual continuous solutions. Though we have not fully justified in this work the spectral decomposition used to derive (3.9), the fact that the infinite dimensional dynamics are so well approximated by the finite dimensional system constructed from these spectral assumptions is quite good evidence that this approximation is a valid one. However, as mentioned in Section 2.1, investigating the validity of spectral assumptions will be an important topic of future research.

## 6. CONCLUSIONS AND DISCUSSION OF FUTURE WORK

In this work, we have used a sinc discretization method to compute the coefficients of the dynamical system (3.9), which is valid near the minimal mass soliton for a saturated nonlinear Schrödinger equation. We find that the dynamical system is an accurate approximation to the full nonlinear solution in a neighborhood of the minimal mass. Moreover, we see that there are two distinct regimes of the dynamical system.

The first regime given by  $\rho_4(0) > 0$  represents oscillation along the soliton curve. The finite dimensional oscillations are valid solutions on long time scales in the conservative PDE, hence we may observe long time closeness of our finite dimensional approximation to the full solution of (1.1).




FIGURE 3. A plot of the solution to the system of ODE's as well as the full solution to (1.1) derived for solutions near the minimal soliton for  $\rho_3(0) > 0$  and  $\rho_3(0) < 0, \rho_4(0) > 0$  for  $\omega_0 = .177588, N = 1000$ .

The second regime given by  $\rho_4(0) < 0$  is the dispersive regime, which results in motion along the unstable branch of the soliton curve. Indeed, a perturbation of this type reduces the mass, which near the minimal mass should lead to dispersion in the primitive system. Obviously, we eventually leave the regime of validity for our approximation. But






FIGURE 4. A plot of the solution to the system of ODE's as well as the full solution to (1.1) derived for solutions near the minimal soliton for  $\rho_3(0) > 0$  and  $\rho_3(0) < 0, \rho_4(0) > 0$  for  $\omega_0 = .17, N = 500$ .

it is notable that on the shorter time scales, our system captures some aspect of the full dynamics.

We cannot numerically verify our conjecture that soliton preserving perturbations of unstable solitons dynamically select stable solitons. However, when we begin with perturbations that are expected to continue to have a soliton component, we see oscillations about




FIGURE 5. A plot of the solution to the system of ODE's as well as the full solution to (1.1) derived for solutions near the minimal soliton for  $\alpha_0 = \rho_3(0) > 0$  and  $\alpha_0 = \rho_3(0) < 0$ ,  $\beta_0 = \rho_4(0) > 0$  for  $\omega_0 = .15$ ,  $N = 500$ .

the minimal mass; this strongly suggests that, through coupling to the continuous spectrum, the oscillations will damp towards a near-minimal-mass stable soliton. This would be quite satisfying from a physical perspective as the system would be moving towards the configuration of lowest energy in some sense.




FIGURE 6. A plot of the solution to the system of ODE's as well as the full solution to (1.1) derived for solutions near the minimal soliton for  $\alpha_0 = \rho_3(0) > 0$  and  $\alpha_0 = \rho_3(0) < 0, \beta_0 = \rho_4(0) < 0$  for  $\omega_0 = .177588, N = 1000$ .

Our ultimate objective is to rigorously characterize the stability properties of the minimal mass soliton. We conjecture that perturbations giving rise to oscillatory dynamics in the finite dimensional system will, in the primitive equation, transition to a stable soliton. Excess mass will be lost by radiation damping. Likely, using current techniques this analysis can only be truly done in a perturbative setting, though we also conjecture that initial




FIGURE 7. A plot of the solution to the system of ODE's as well as the full solution to (1.1) derived for solutions near the minimal soliton for  $\alpha_0 = \rho_3(0) > 0$  and  $\alpha_0 = \rho_3(0) < 0, \beta_0 = \rho_4(0) < 0$  for  $\omega_0 = .17, N = 500$ .

conditions near strongly unstable solitons should exhibit similar behavior. Hopefully more powerful techniques will eventually be developed for the global study of the stable soliton curve as an attractor of the full nonlinear dynamics.




FIGURE 8. A plot of the solution to the system of ODE's as well as the full solution to (1.1) derived for solutions near the minimal soliton for  $\alpha_0 = \rho_3(0) > 0$  and  $\alpha_0 = \rho_3(0) < 0, \beta_0 = \rho_4(0) < 0$  for  $\omega_0 = .15, N = 500$ .

## REFERENCES

- [1] M.J. Ablowitz and Z.H. Musslimani. Spectral renormalization method for computing self-localized solutions to nonlinear systems. *Optics letters*, 30(16):2140–2142, 2005.

- [2] G. Akrivis, V. Dougalis, O. Karakashian, and W. McKinney. On fully discrete Galerkin methods of second-order temporal accuracy for the nonlinear Schrödinger equation. *Numerische Mathematik*, 59(194):31–53, 1991.
- [3] G. Akrivis, V. Dougalis, O. Karakashian, and W. McKinney. Solving the systems of equations arising in the discretization of some nonlinear p.d.e's by implicit Runge-Kutta methods. *RAIRO Modl. Math. Anal. Numr.*, 31(3):251–288, 1997.
- [4] G. Akrivis, V. Dougalis, O. Karakashian, and W. McKinney. Numerical approximation of blow-up of radially symmetric solutions of the nonlinear Schrödinger equation. *SIAM Journal of Scientific Computing*, 25(1):186–212, 2003.
- [5] K. Al-Khaled. Sinc numerical solution for solitons and solitary waves. *Journal of Computational and Applied Mathematics*, 130(1-2):283–292, 2001.
- [6] E.L. Allgower and K. Georg. *Numerical Continuation Methods: An Introduction*. Springer, 1990.
- [7] W. Bao and Q. Du. Computing the ground state solution of Bose-Einstein condensates by a normalized gradient flow. *SIAM Journal of Scientific Computing*, 25:1674–1697, 2004.
- [8] I.V. Barashenkov, A.D. Gocheva, V.G. Makhankov, and I.V. Puzynin. Stability of the soliton-like “bubbles”. *Physica D*, 34:240–254, 1989.
- [9] I.V. Barashenkov and E.Y. Panova. Stability and evolution of the quiescent and travelling solitonic bubbles. *Physica D*, 69:114–134, 1993.
- [10] N. Bellomo. Nonlinear models and problems in applied sciences from differential quadrature to generalized collocation methods. *Mathematical and Computer Modelling*, 26(4):13–34, 1997.
- [11] N. Bellomo and L. Ridolfi. Solution of nonlinear initial-boundary value problems by sinc collocation-interpolation methods. *Computers and Mathematics with Applications*, 29(4):15–28, 1995.
- [12] H. Berestycki and P.L. Lion. Nonlinear scalar field equations, I: Existence of a ground state. *Arch. Rational. Mech. anal.*, 82(4):313–345, 1983.
- [13] B. Bialecki. Sinc-type approximations in H 1-norm with applications to boundary value problems. *Journal of Computational and Applied Mathematics*, 25(3):289–303, 1989.
- [14] B. Bialecki. Sinc-collocation methods for two-point boundary value problems. *IMA Journal of Numerical Analysis*, 11(3):357–375, 1991.
- [15] J.P. Boyd. *Chebyshev and Fourier Spectral Methods*. Courier Dover Publications, 2001.
- [16] V. Buslaev and V. Grikurov. Simulation of Instability of Bright Solitons for NLS with Saturating Nonlinearity. *Math. Comput. Simulation*, pages 539–546, 2001.
- [17] V.S. Buslaev and G.S. Perelman. On the stability of solitary waves for nonlinear Schrödinger equations. *Amer. Math. Soc. Transl. Ser. 2*, 164(1):75–98, 1995.
- [18] T.S. Carlson, J. Dockery, and J. Lund. A sinc-collocation method for initial value problems. *Mathematics of Computation*, 66(217):215–235, 1997.
- [19] T. Cazenave. *Semilinear Schrodinger Equations*, volume 10 of *Courant Lecture Notes in Mathematics*. American Mathematical Society, 2003.
- [20] A. Comech and D. Pelinovsky. Purely Nonlinear Instability of Standing Waves with Minimal Energy. *Communications on Pure and Applied Mathematics*, 56:1565–1607, 2003.
- [21] Cuccagna. On asymptotic stability of ground states of NLS. *Rev. Math. Phys.*, 15(8):877–903, 2003.
- [22] L. Demanet and W. Schlag. Numerical verification of a gap condition for a linearized nonlinear Schrödinger equation. *Nonlinearity*, pages 829–852, 2006.
- [23] K. Dimitrevski, E. Reimhult, E. Svensson, A. Ohgren, D. Anderson, A. Berntson, M. Lisak, and M.L. Quiroga-Teixeiro. Analysis of stable self-trapping of laser beams in cubic-quintic nonlinear media. *Physics Letters A*, 248:369–376, 1998.
- [24] A. Eisner and B. Turkington. Nonequilibrium statistical behavior of nonlinear Schrödinger equations. *Physica D*, 213:85–97, 2006.

- [25] M. El-Gamel. Sinc and the numerical solution of fifth-order boundary value problems. *Applied Mathematics and Computation*, 187(2):1417–1433, 2007.
- [26] M. El-Gamel, S.H. Behiry, and H. Hashish. Numerical method for the solution of special nonlinear fourth-order boundary value problems. *Applied Mathematics and Computation*, 145(2-3):717–734, 2003.
- [27] M. El-Gamel and AI Zayed. Sinc-Galerkin method for solving nonlinear boundary-value problems. *Computers and Mathematics with Applications*, 48(9):1285–1298, 2004.
- [28] B. Erdogan and W. Schlag. Dispersive estimates for Schrödinger operators in the presence of a resonance and/or eigenvalue at zero energy in dimension three. II. *J. Anal. Math*, 99:199–248, 2006.
- [29] M. Grillakis, J. Shatah, and W. Strauss. Stability theory of solitary waves in the presence of symmetry. ii. *J. Funct. Anal.*, 94(2):308–348, 1990.
- [30] J. Holmer, J. Marzuola, and M. Zworski. Soliton splitting by external delta potentials. *Journal of Nonlinear Science*, 17(4):349–367, 2007.
- [31] T.W. Johnston, F. Vidal, and D. Fréchette. Laser-plasma filamentation and the spatially periodic nonlinear Schrödinger equation approximation. *Physics of Plasmas*, 4(5):1582–1588, 1997.
- [32] Eric Jones, Travis Oliphant, Pearu Peterson, et al. SciPy: Open source scientific tools for Python, 2001–.
- [33] C. Josserand, Y. Pomeau, and S. Rica. Cavitation versus vortex nucleation in a superfluid model. *Physical Review Letters*, 75(17):3150–3153, 1995.
- [34] C. Josserand and S. Rica. Coalescence and droplets in the subcritical nonlinear Schrödinger equation. *Physical Review Letters*, 78(7):1215–1218, 1997.
- [35] J. Krieger and W. Schlag. Stable manifolds for all monic supercritical focusing nonlinear Schrödinger equations in one dimension. *J. Amer. Math. Soc.*, 19(4):815–920, 2006.
- [36] T.I. Lakoba and J. Yang. A generalized Petviashvili iteration method for scalar and vector Hamiltonian equations with arbitrary form of nonlinearity. *Journal of Computational Physics*, 226(2):1668–1692, 2007.
- [37] B. J. LeMesurier, G. Papanicolaou, C. Sulem, and P.-L. Sulem. Focusing and multi-focusing solutions of the nonlinear Schrödinger equation. *Phys. D*, 31(1):78–102, 1988.
- [38] J. Lund and K.L. Bowers. *Sinc Methods for Quadrature and Differential Equations*. Society for Industrial Mathematics, 1992.
- [39] L. Lundin. A Cardinal Function Method of Solution of the Equation  $\Delta u = u - u^3$ . *Mathematics of Computation*, 35(151):747–756, 1980.
- [40] J. Marzuola and G. Simpson. Numerical and analytic results on the spectrum of matrix Hamiltonian operators. In preparation, 2009.
- [41] K. McCleod. Uniqueness of Positive Radial Solutions of  $\Delta u + f(u) = 0$  in  $\mathbb{R}^n$ , II. *Transactions of the American Mathematical Society*, 339(2):495–505, 1993.
- [42] A. Mohsen and M. El-Gamel. On the Galerkin and collocation methods for two-point boundary value problems using sinc bases. *Computers and Mathematics with Applications*, 56(4):930–941, 2008.
- [43] D.E. Pelinovsky, V.V. Afanasjev, and Y.S. Kivshar. Nonlinear theory of oscillating, decaying, and collapsing solitons in the generalized nonlinear Schrödinger equation. *Physical Review E*, 53(2):1940–1953, 1996.
- [44] V.I. Petviashvili. Equation of an extraordinary soliton. *Soviet Journal of Plasma Physics*, 2:469–472, 1976.
- [45] M. Quiroga-Teixeiro and H. Michinel. Stable azimuthal stationary state in quintic nonlinear optical media. *Journal of the Optical Society of America B*, 14(8):2004–2009, 1997.
- [46] R. Revelli and L. Ridolfi. Sinc collocation-interpolation method for the simulation of nonlinear waves. *Computers and Mathematics with Applications*, 46(8-9):1443–1453, 2003.

- [47] I. Rodnianski, W. Schlag, and A. Soffer. Asymptotic stability of  $N$ -soliton states of NLS. Preprint, 2003.
- [48] W. Schlag. Stable manifolds for an orbitally unstable NLS. To appear in *Annals of Math*, 2004.
- [49] J. Shatah. Stable Standing Waves of Nonlinear Klein-Gordon Equations. *Communications in Mathematical Physics*, 91:313–327, 1983.
- [50] J. Shatah. Unstable Ground State of Nonlinear Klein-Gordon Equations. *Transactions of the American Mathematical Society*, 290(2):701–710, 1985.
- [51] J. Shatah and W. Strauss. Instability of Nonlinear Bound States. *Communications in Mathematical Physics*, 100:173–190, 1985.
- [52] G. Simpson and M Spiegelman. Robust numerical benchmarks for magma dynamics. In preparation.
- [53] A. Soffer and M.I. Weinstein. Resonances, radiation damping and instability in Hamiltonian nonlinear wave equations. *Invent. Math.*, 136(1):9–74, 1999.
- [54] F. Stenger. A” Sinc-Galerkin” method of solution of boundary value problems. *Mathematics of Computation*, pages 85–109, 1979.
- [55] F. Stenger. Numerical methods based on the whittaker cardinal, or sinc functions. *SIAM Review*, 23:165–224, 1981.
- [56] F. Stenger. *Numerical Methods Based on Sinc and Analytic Functions*. Springer, 1993.
- [57] F. Stenger. Summary of sinc numerical methods. *Journal of Computational and Applied Mathematics*, 121(1-2):379–420, 2000.
- [58] C. Sulem and P. Sulem. *The Nonlinear Schrödinger Equation. Self-focusing and wave-collapse*, volume 39 of *Applied Mathematical Sciences*. Springer-Verlag, 1999.
- [59] V. Tikhonenko, J. Christou, and B. Luther-Davies. Three dimensional bright spatial soliton collision and fusion in a saturable nonlinear medium. *Physical Review Letters*, 76(15):2698–2701, 1996.
- [60] M. I. Weinstein. Modulational stability of ground states of nonlinear Schrödinger equations. *SIAM J. Math. Anal.*, 16:472–491, 1985.
- [61] M. I. Weinstein. Lyapunov stability of ground states of nonlinear dispersive evolution equations. *Comm. Pure Appl. Math.*, 39:472–491, 1986.
- [62] E.M. Wright, B.L. Lawrence, W. Torruellas, and G. Stegeman. Stable self-trapping and ring formation in polydiacetylene para-toluene sulfonate. *Optics Letters*, 20(24):2481, 1995.
- [63] J. Yang and T.I. Lakoba. Universally-convergent squared-operator iteration methods for solitary waves in general nonlinear wave equations. *Studies in Applied Mathematics*, 118(2):153, 2007.
- [64] J. Yang and T.I. Lakoba. Accelerated imaginary-time evolution methods for the computation of solitary waves. *Studies in Applied Mathematics*, 120(3):265–292, 2008.

## APPENDIX A. DETAILS OF NUMERICAL METHODS

**A.1. Sinc Approximation.** Here we briefly review sinc and its properties. The texts [38, 56] and the articles [55, 10, 57] provide an excellent overview. As noted, sinc collocation and Galerkin schemes have been used to solve a variety of partial differential equations.

Recall the definition of sinc,

$$(A.1) \quad \text{sinc}(z) \equiv \begin{cases} \frac{\sin(\pi z)}{\pi z}, & \text{if } z \neq 0 \\ 1, & \text{if } z = 0 \end{cases}.$$



and for any  $k \in \mathbb{Z}$ ,  $h > 0$ , let

$$(A.2) \quad S(k, h)(x) = \operatorname{sinc} \left( \frac{x - kh}{h} \right).$$

The sinc function can be used to exactly represent functions in the Paley-Wiener class. We spectrally represent functions with sinc in a weaker function space. First, we define a *strip* in the complex plane,

$$(A.3) \quad D_d = \{z \in \mathbb{C} \mid |\operatorname{Im} z| < d\}.$$

Then we define the function space:

**Definition A.1.**  $B^p(D_d)$  is the set of analytic functions on  $D_d$  satisfying:

$$(A.4a) \quad \|f(t + i\cdot)\|_{L^1(-d, d)} = O(|t|^a), \quad \text{as } t \rightarrow \pm\infty, \text{ with } a \in [0, 1),$$

$$(A.4b) \quad \lim_{y \rightarrow d^-} \|f(\cdot + iy)\|_{L^p} + \lim_{y \rightarrow -d^-} \|f(\cdot - iy)\|_{L^p} < \infty.$$

Then, we have the following

**Theorem 1.** (Theorem 2.16 of [38])

Assume  $f \in B^p(D_d)$ ,  $p = 1$  or  $2$ , and  $f$  satisfies the decay estimate

$$(A.5) \quad |f(x)| \leq C e^{-\alpha|x|}.$$

If  $h$  is selected such that

$$(A.6) \quad h = \sqrt{\pi d / (\alpha M)} \leq \min \left\{ \pi d, \pi / \sqrt{2} \right\},$$

then

$$\|\partial_x^n f - \partial_x^n C_M(f, h)\|_{L^\infty} \leq C M^{(n+1)/2} e^{(-\sqrt{\pi d \alpha M})}.$$

$d$  identifies a strip in the complex plane, of width  $2d$ , about the real axis in which  $f$  is analytic. This parameter may not be obvious; others have found  $d = \pi/2$  sufficient.

For the NLS equation of order  $2\sigma + 1$ ,

$$d_{\text{NLS}} = \frac{\pi}{\sqrt{2\omega\sigma}}.$$

Saturated NLS “interpolates” between second and seventh order NLS. We thus reason it is fair to take  $d = \pi/\sqrt{6\omega}$ . Though we do not prove that the soliton and the associated elements of the kernel lie in these  $B^p(D_d)$  spaces or satisfy the hypotheses of Theorem 1, we use (A.6) to guide our selection of an optimal  $h$ . Since the soliton has  $\alpha = \sqrt{\omega}$ , we reason that it should be acceptable to take

$$(A.7) \quad h = \sqrt{\frac{\pi d}{\alpha M}} = \sqrt{\frac{\pi^2}{6\omega M}}.$$

(A.7) is dependent on both  $M$  and  $\omega$ . Were we to use (A.7) as is, it would complicate approximating, amongst other things, the derivative with respect to  $\omega$  of the soliton. To

avoid this, we use *a priori* estimates on  $\omega^*$ , given in Appendix A.4.1. Since we know that  $\omega^* \sim .18 < .25$ , it is sufficient to take

$$d = \pi\sqrt{2/3}.$$

Likewise, since  $\omega^* > .1$ , we may take

$$\alpha = \sqrt{1/10}.$$

Thus, instead of (A.7), we use

$$(A.8) \quad h = \pi\sqrt{\frac{\sqrt{20/3}}{M}}.$$

We conjecture that this is a valid grid spacing for all  $\omega \in (.1, .25)$ ; our computations are consistent with this assumption.

**A.2. Numerical Continuation.** As discussed in Section 4.1, the discrete system approximating (1.5) is

$$(A.9) \quad \vec{F}(\vec{\phi}) = D^{(2)}\vec{\phi} - \omega\vec{\phi} + g(\vec{\phi})\vec{\phi} = 0.$$

The multiplication in  $g(\vec{\phi})\vec{\phi}$  is performed elementwise. In order to solve this discrete system, we need a good starting point for our nonlinear solver. We produce this guess by numerical continuation.

Define the function

$$\hat{g}(x; \tau) = \frac{x^3}{1 + \tau x^2}.$$

Note that  $\hat{g}(x, 0)$  is 7-th order NLS and  $\hat{g}(x, 1) = g(x)$ , saturated NLS. We now solve

$$(A.10) \quad \vec{G}(\vec{\phi}; \tau) = D^{(2)}\vec{\phi} - \lambda\vec{\phi} + \hat{g}(\vec{\phi}; \tau)\vec{\phi} = 0.$$

At  $\tau = 0$ , the analytic NLS soliton serves as the initial guess for computing  $\vec{\phi}_{\tau=0}$ .  $\vec{\phi}_{\tau=0}$  is then the initial guess for solving (A.10) at  $\tau = \Delta\tau$ . We iterate in  $\tau$  until we reach  $\tau = 1$ . This is numerical continuation in the artificial parameter  $\tau$ , [6]. This process succeeds with relatively few steps of  $\Delta\tau$ ; in fact only  $O(10)$  steps are required.

**A.3. Convergence Data.** Table 2 offers some examples of the robust and rapid convergence seen in the coefficients of (3.9). These values are all computed at the minimal mass soliton. Also see Table 1.

**A.4. Comparisons with Other Methods.** We can benchmark our sinc algorithm against several other methods. Available algorithms include numerical quadrature along with more recent approaches such as spectral renormalization, also called the Petviashvili method [44, 1, 36], the imaginary time method [7, 64], and the squared operator method [63].

TABLE 2. The convergence of several coefficients for the ODE system, computed at  $\omega^*$ .

$M$	$a_0$	$c_{14}$	$p_{13}$	$n_{133}$
20	-0.54851448504	6.04829942099	-6.04829942099	8.79376331231
40	-0.553577138662	6.12521927361	-6.12521927361	8.81625889017
60	-0.553555163933	6.12811479039	-6.12811479039	8.81709463059
80	-0.553550441653	6.12827391288	-6.12827391288	8.81713847275
100	-0.553549989603	6.12828576495	-6.12828576495	8.81714173314
200	-0.553549934797	6.12828700415	-6.12828700415	8.81714206225
300	-0.553549934793	6.12828700423	-6.12828700423	8.81714206227
400	-0.553549934795	6.12828700421	-6.12828700421	8.81714206223
500	-0.553549934794	6.12828700423	-6.12828700423	8.81714206227

TABLE 3. Value of the coefficients in (3.9) computed with  $M = 200$ .

Coefficient	Value
$g_{33}$	-6.61999411752
$g_{44}$	-12.4582451458
$c_{14}$	6.12828700415
$c_{23}$	1.46358108488
$c_{34}$	4.0422919871
$c_{43}$	0.131304385722
$p_{13}$	-6.12828700415
$p_{24}$	-17.9305799071
$p_{33}$	6.61999411752
$p_{44}$	12.4582451458
$n_{133}$	8.81714206225
$n_{144}$	1.84068246508
$n_{234}$	1.45559877602
$n_{333}$	-0.792198288158
$n_{344}$	0.013887281387
$n_{434}$	-0.0822482271619
$a_0$	-0.553549934797

A.4.1. *Quadrature Methods.* The soliton equation may be integrated once to get

$$(A.11) \quad \frac{1}{2}(\partial_x \phi)^2 - \frac{1}{2}\lambda\phi^2 + \frac{1}{4}[\phi^4 - \log(1 + \phi^4)] = 0.$$

Equation (A.11) yields an implicit algebraic expression for the amplitude,  $\phi(0)$ ,

$$(A.12) \quad -\frac{1}{2}\lambda\phi(0)^2 + \frac{1}{4}[\phi(0)^4 - \log(1 + \phi(0)^4)] = 0.$$

Using (A.11) and (A.12), we can express the mass as

$$\|\phi\|_{L^2}^2 = \int_{-\infty}^{\infty} \phi(x)^2 dx = 2 \int_0^{\infty} \phi(x)^2 dx = 2 \int_0^{\phi(0)} \rho^2 \left\{ \lambda \rho^2 - \frac{1}{2} [\rho^4 - \log(1 + \rho^4)] \right\}^{-1/2} d\rho.$$

Thus, the mass of the soliton with parameter  $\omega$  is

$$(A.13) \quad \|\phi_\omega\|_{L^2}^2 = 2 \int_0^{\phi(0;\omega)} \rho^2 \left\{ \omega \rho^2 - \frac{1}{2} [\rho^4 - \log(1 + \rho^4)] \right\}^{-1/2} d\rho.$$

Equations (A.12) and (A.13) can be used to approximate  $\omega^*$  by numerically minimizing (A.13). To compute the amplitude of the soliton, we solve (A.12) using Brent's method with a tolerance of 1.0e-14. We use the singular integral integrator QAGS from QUADPACK, which for this problem is, unfortunately, limited to a relative error of 5.0e-12 and an absolute error of 1.0e-15. Trying different routines from the optimization module of SciPy, [32], we summarize our results in Table 4, which contains data from our sinc computations. There is a spread of  $O(1e-12)$  amongst the computed minimal masses and a spread of  $O(1e-7)$  amongst the  $\omega^*$ . These differences are consistent with the prescribed relative error of the quadrature, suggesting the precision of this approach to computing the minimal mass and associated  $\omega$  is limited by the quadrature algorithm. Note that we do not compute the solitary wave with this technique; we merely identify the minimal mass soliton parameter and the mass of that soliton.

*A.4.2. Spectral Renormalization Methods.* A Fourier transform may be applied to the soliton equation to get

$$(A.14) \quad \hat{\phi}_\omega(k) = \frac{\mathcal{F}(g(\phi_\omega^2)\phi_\omega)(k)}{k^2 + \omega}$$

where  $k$  is the wave number. Let us introduce variable  $w(x)$ , with  $\phi_\omega(x) = \theta w(x)$ , where  $\theta$  is an unknown, nonzero, constant. Introducing this into (A.14), we have

$$(A.15) \quad \hat{w}(k) = \frac{\mathcal{F}(g(\theta^2 w^2)w)(k)}{k^2 + \omega} \equiv \mathcal{Q}_\theta[\hat{w}](k).$$

Multiplying (A.15) by  $\hat{w}(k)^*$ , the complex conjugate, and integrating over all  $k$ , we compute

$$(A.16) \quad G(\theta; \hat{w}) \equiv \int |\hat{w}(k)|^2 dk - \int \hat{w}(k)^* \mathcal{Q}_\theta[\hat{w}](k) dk = 0.$$

This may be interpreted as a constraint on  $\theta$ . This motivates the iteration described in [1]. Suppose we know  $w_m(x)$  and  $\theta_m$ , a pair of approximations of the true values. To get the next approximation, we compute

$$(A.17) \quad \hat{w}_{m+1}(k) = \mathcal{Q}_{\theta_m}[\hat{w}_m](k)$$

and then solve

$$(A.18) \quad G(\theta; \hat{w}_{m+1}) = 0$$

TABLE 4. The soliton parameter and mass of the minimal mass soliton computed by both quadrature and spectral renormalization. Also included is some of the data for the sinc method appearing in Table 1.

Algorithm	$\omega^*$	$\int  \phi ^2 dx$
fminbound	0.177588368745261	3.821149022780204
Brent	0.177587963826864	3.821149022778472
golden	0.177587925853761	3.821149022776717
Spec. Re.	0.177588064106709	3.821149022780361
sinc with $M = 100$	0.177588063805561	3.821149022493814
sinc with $M = 200$	0.177588065432740	3.821149022780618
sinc with $M = 400$	0.177588065433095	3.821149022780896

for  $\theta_{m+1}$ . We repeat this until the sequence  $\{w_m\}$  satisfies our convergence criteria. From this we then recover  $\phi_\omega$ . The advantage of this is it can use the fast Fourier transform. Using spectral renormalization, we then minimize the approximate mass numerically. This is readily implemented in MATLAB, using `fzero` to solve (A.18) for a given value of  $\omega$  and `fminbnd` to find the minimal mass value of  $\omega$ . We iterate in  $w$  until either  $\|w_{m+1} - w_m\|_{\ell^2} < \text{Abs. Tol.}$  or  $\|w_{m+1} - w_m\|_{\ell^2} / \|w_{m+1}\|_{\ell^2} < \text{Rel. Tol.}$ . This is performed with a relative and absolute tolerances of  $1e-15$  in the spectral renormalization component, and a tolerance of  $1e-15$  in both `fminbnd` `fzero`. Our spatial domain is  $[-200, 200)$  with  $2^{12}$  grid points. As seen in Table 4, this is in good agreement with both the sinc method and the quadrature method.

*E-mail address:* `jm3058@columbia.edu`

DEPARTMENT OF APPLIED MATHEMATICS, COLUMBIA UNIVERSITY, 200 S. W. MUDD, 500 W. 120TH ST., NEW YORK CITY, NY 10027, USA

*E-mail address:* `raynorsg@math.wfu.edu`

MATHEMATICS DEPARTMENT, WAKE FOREST UNIVERSITY, P.O. BOX 7388, 127 MANCHESTER HALL, WINSTON-SALEM, NC 27109, USA

*E-mail address:* `simpson@math.toronto.edu`

MATHEMATICS DEPARTMENT, UNIVERSITY OF TORONTO, 40 ST. GEORGE ST., TORONTO, ONTARIO, CANADA M5S 2E4