

Climate-independent paleoaltimetry using stomatal density in fossil leaves as a proxy for CO₂ partial pressure: Comment and Reply

COMMENT

Daniel M. Johnson
William K. Smith
Miles R. Silman

Department of Biology, Wake Forest University, Winston-Salem, North Carolina 27109-7325, USA

Stomatal density (SD) has been shown to change with CO₂ concentration (at the same atmospheric pressure; Lake et al., 2001), and SD of fossilized leaf material or leaf imprints can be used to indicate past CO₂ concentrations (Beerling et al., 1998). However, the results of the analytical approach described by McElwain (2004) do not include an important parameter influencing the effect of altitude on CO₂ diffusion and uptake by a plant. Although the partial pressure of carbon dioxide does decrease in a predictable manner with decreasing atmospheric pressure, the diffusion of all gas-phase molecules increases at lower atmospheric pressure (Fig. 1). As a result, photosynthetic CO₂ uptake does not appear to become more rate-limited at greater altitudes, due primarily to the compensatory effects of increased diffusion rates (e.g., Smith and Donahue, 1991; Terashima et al., 1995). Thus, there would be no selective force for increasing SD in plant species evolving at higher altitudes, and the relationship suggested by McElwain (2004) is not expected to be generally applicable.

Analytically, diffusion is often expressed according to Fick's Law as

$$\text{Flux density of } j \text{ (mol m}^{-2} \text{ s}^{-1}) = D_j (\Delta C_j) / \Delta x, \quad (1)$$

where D_j is the diffusion coefficient for the molecule in question, ΔC_j is the gradient in the concentration or partial pressure of molecule j , and Δx is the diffusing distance (Nobel, 1999). As seen in Figure 1, both D_j and ΔC_j influence flux density proportionally. Moreover, both terms change in similar proportions with increasing altitude, but are inversely related. For example, an increase in altitude from sea level to 4 km results in a decrease in the partial pressure of CO₂ of -38% (0.37–0.23 kPa), and an increase in D_c of 53% (1.65–2.53 m² s⁻¹ × 10⁻⁵) based on a relatively dry adiabatic lapse rate of 8 °C per 1000 m (Fig. 1).

A host of abiotic factors may be associated with changes in altitude, most of which can strongly influence stomatal density (e.g., Tichá, 1982; Lockheart et al., 1998; Qiang et al., 2003). Numerous studies have found increases in SD with elevation (e.g., Hovenden and Brodribb, 2000), decreases (e.g., Hultine and Marshall, 2000), or have been inconclusive (e.g., Qiang et al., 2003), suggesting that CO₂ concentration is not a primary altitudinal factor driving SD.

Although SD has been used to estimate historical CO₂ concentrations on a geological time scale, the similar proportional changes in D_{CO_2} and CO₂ partial pressure would indicate a compensating influence of the two acting in concert. As mentioned above, this compensating relationship has already been addressed in other publications dealing with the impact of altitude on the photosynthetic process. Similarly, there is no convincing evidence that greater altitude results in greater SD as a general rule. In this regard, it is actually total pore area per unit leaf area (which incorporates both stomatal size and frequency) that is the most pertinent parameter, and one which is rarely measured (e.g., Smith and Knapp, 1990). However, others have also concluded that the greatest impact of altitude is probably on plant transpiration, primarily because the increasing dryness of the air with altitude is often exacerbated by the concomitant increase in the diffusion coefficient of water vapor in air (D_{w}) (Gale, 1972; Smith and Geller, 1979; Smith and Knapp, 1990; Terashima et al., 1995).




Figure 1. A: Change in CO₂ partial pressure ($p\text{CO}_2$). B: The diffusion coefficient for CO₂ in air (D_{CO_2}) with altitude. Curves represent absolute values (closed circles) and percent changes from sea-level values (open circles). Curve for $D_{\text{CO}_2} = 1.652 \times 10^{-5} (T/273)^{1.8} (P_0/P)$ (Nobel, 1999).

REFERENCES CITED

- Beerling, D.J., McElwain, J.C., and Osborne, C.P., 1998, Stomatal responses of the 'living fossil' *Ginkgo biloba* L. to changes in atmospheric CO₂ concentrations: *Journal of Experimental Botany*, v. 49, p. 1603–1607, doi: 10.1093/jexbot/49.326.1603.
- Gale, J., 1972, Elevation and transpiration: Some theoretical considerations with special reference to mediterranean-type climate: *Journal of Applied Ecology*, v. 9, p. 691–701.
- Hovenden, M.J., and Brodribb, T., 2000, Altitude of origin influences stomatal conductance and therefore maximum assimilation rate in Southern Beech, *Nothofagus cunninghamii*: *Australian Journal of Plant Physiology*, v. 27, p. 451–456.
- Hultine, K.R., and Marshall, J.D., 2000, Altitude trends in conifer leaf morphology and stable isotope composition: *Oecologia*, v. 123, p. 32–40, doi: 10.1007/s004420050986.
- Lake, J.A., Quick, W.P., Beerling, D.J., and Woodward, F.I., 2001, Signals from mature to new leaves: *Nature*, v. 411, p. 154, doi: 10.1038/35075660.
- Lockheart, M.J., Poole, I., Van Bergen, P.F., and Evershed, R.P., 1998, Leaf carbon isotope compositions and stomatal characters: Important considerations for paleoclimate reconstructions: *Organic Geochemistry*, v. 29, p. 1003–1008, doi: 10.1016/S0146-6380(98)00168-5.
- McElwain, J.C., 2004, Climate-independent paleoaltimetry using stomatal density in fossil leaves as a proxy for CO₂ partial pressure: *Geology*, v. 32, p. 1017–1020, doi: 10.1130/G20915.1.
- Nobel, P., 1999, *Physiochemical and environmental physiology*: San Diego, Academic Press, 424 p.

- Qiang, W.Y., Wang, X.L., Chen, T., Feng, H.Y., An, L.Z., He, Y.Q., and Wang, G., 2003, Variations in stomatal density and carbon isotope values of *Picea crassifolia* at different altitudes in the Qilian Mountains: *Trees*, v. 17, p. 258–262.
- Smith, W.K., and Donahue, R., 1991, Simulated effect of altitude on photosynthetic CO₂ uptake potential in plants: *Plant, Cell & Environment*, v. 14, p. 133–136.
- Smith, W.K., and Geller, G., 1979, Plant transpiration at high elevations: Theory, field measurement, and comparisons with desert plants: *Oecologia*, v. 41, p. 109–122, doi: 10.1007/BF00344841.
- Smith, W.K., and Knapp, A.K., 1990, Ecophysiology of high elevation forests, in Osmond, C.B., et al., eds., *Plant Biology of the Basin and Range*: New York, Springer-Verlag, p. 87–142.
- Terashima, I., Masuzawa, T., Ohba, H., and Yokoi, Y., 1995, Is photosynthesis suppressed at higher elevation due to low CO₂ pressure?: *Ecology*, v. 76, p. 2663–2668.
- Tichá, I., 1982, Photosynthetic characteristics during ontogenesis of leaves: 7. Stomata density and sizes: *Photosynthetica*, v. 16, p. 375–471.

REPLY

Jennifer C. McElwain

Department of Geology, The Field Museum, Chicago, Illinois 60605-2496

The functional explanation for increased plant leaf stomatal density (SD) with elevation remains a topic of much debate among plant ecophysiologicalists (Friend and Woodward, 1990; Gale, 1972, 2004; Smith and Donahue, 1991; Terashima et al., 1995; Woodward and Bazzaz, 1988). Johnson et al. raise an interesting aspect of this ongoing debate. Based on the observation that gaseous diffusion increases with decreasing atmospheric pressure these authors argue that any negative impact of decreasing carbon dioxide partial pressure ($p\text{CO}_2$) with elevation on plant photosynthesis would likely be compensated for by greater diffusion rates of CO₂ into the leaf. Using these theoretical considerations Johnson et al. argue that decreasing CO₂ partial pressure is unlikely to select for increased SD in plant species evolving at high elevation, and thus, the highly significant relationship demonstrated between Californian black oak SD and elevation is unlikely to be generally applicable. In their theoretical calculations, however, Johnson et al. fail to take into account the observation that transpiration rates are often extremely elevated at high elevation due to (1) higher light intensity, (2) increased diffusion of H₂O in air at reduced atmospheric pressure, and (3) an increased density gradient of H₂O vapor out of the leaf into the ambient air (Gale, 2004). As H₂O vapor efflux through stomata has been shown to hamper CO₂ diffusion into the leaf (von Caemmerer and Farquhar, 1981), it is highly likely that increased transpirational water vapor loss at higher elevations would act to cancel out the higher diffusion coefficient of CO₂ at reduced barometric pressure (Gale, 2004), thereby selecting strongly for increased SD at high elevation to maintain adequate conductance to CO₂ diffusion.

Johnson et al. argue that many abiotic factors in addition to $p\text{CO}_2$ change with altitude and suggest that these other factors rather than $p\text{CO}_2$ may be responsible for the observed increase in SD in *Quercus kelloggii*. The potential role of other biotic and abiotic factors on SD, including temperature, light intensity, water availability, sex of the plant, etc., is well known (McElwain and Chaloner, 1996; Royer, 2001). Carbon dioxide and to a much lesser extent light intensity are the only factors which are known to control the actual development of stomata from epidermal initials. The effect of CO₂ on leaf development can be tracked by measuring stomatal index (SI); a ratio of the number of stomata to the total number of cells (stomata plus epidermal) on the leaf surface. In contrast all other biotic/abiotic variables which influence SD do so indirectly (Beerling and Kelly, 1997). In other words, although factors like temperature, water availability, and humidity do not directly influence the absolute number of stomata developing from initials, these variables can influence the size and/or spacing of epidermal cells, resulting in stomata being packed more closely together or further apart resulting in higher or lower densities but importantly the same SI.

The significant increase in both SI and SD with elevation in both sun and shade leaves of Californian black oak which I document in my paper

can only therefore be explained by the effect of increasing CO₂ partial pressure on stomatal development from epidermal initials. Increasing SI rules out the effect of all other biotic and abiotic variables. In addition, to reiterate the results in the paper, a historical investigation into the effect of four climatic variables showed no significant correlation with black oak SD. However, the same historical black oak SD data set showed a highly significant inverse correlation with CO₂ concentration. Contrary to the assertions of Johnson et al. therefore, these two lines of evidence suggest that decreasing CO₂ partial pressure is indeed a primary altitudinal factor driving SD changes in Californian black oak.

This does not simply imply however, and I do not state as such in my paper, that the stomatal $p\text{CO}_2$ paleoaltimeter will be universally applicable. Certain taxa are not CO₂ sensitive—they do not show the classic inverse relationship between $p\text{CO}_2$ and SD or SI. These include many grasses and forbs (Reid et al., 2003) and plants with C4 photosynthesis (Raven and Ramden, 1989). These taxa have evolved alternative strategies for optimizing CO₂ uptake for photosynthesis against transpirational water loss, which negates the need to adjust SD. These include fine-tuned stomatal control which enable much more rapid stomatal opening and closing times within minutes rather than hours (as in grasses) and biochemical means of concentrating CO₂ internally thereby ensuring a high CO₂ gradient from the air to the leaf (as in C4 plants). It is imperative therefore that the stomatal $p\text{CO}_2$ paleoaltimeter is only applied to extant taxa, which are known to be CO₂ sensitive. Meta analysis of woody plant SD and SI responses to historical increases in $p\text{CO}_2$ over the past 200 yr and at naturally elevated CO₂ springs has shown that >70% (n = 44) of the taxa analyzed demonstrate the expected inverse relationship (Royer, 2001). This high CO₂ sensitivity among the majority of woody C3 taxa indicates that increased SD with elevation is likely to be the general rule for this group of plants. Coupled with the fact that woody C3 plants are the most well represented group in the plant fossil record, they are an ideal group for application of the stomatal $p\text{CO}_2$ paleoaltimeter.

REFERENCES CITED

- Beerling, D.J., and Kelly, C.K., 1997, Stomatal density responses of temperate woodland plants over the past seven decades of CO₂ increase: A comparison of Salisbury (1927) with contemporary data: *American Journal of Botany*, v. 84, p. 1572–1583.
- Friend, A.D., and Woodward, F.I., 1990, Evolutionary and ecophysiological responses of mountain plants to the growing season environment: *Advances in Ecological Research*, v. 20, p. 59–124.
- Gale, J., 1972, Availability of carbon dioxide at high altitudes: Theoretical considerations: *Ecology*, v. 53, p. 494–497.
- Gale, J., 2004, Plants and altitude—Revisited: *Annals of Botany*, v. 94, p. 199, doi: 10.1093/aob/mch143.
- McElwain, J.C., and Chaloner, W.G., 1996, The fossil cuticle as a skeletal record of environmental change: *Palaios*, v. 11, p. 376–388.
- Raven, J.A., and Ramden, H.J., 1989, Similarity of stomatal index in the C4 plant *Salsola kali* L. in material collected in 1943 and in 1987: Relevance to changes in atmospheric CO₂ content: *Transactions of the Botanical Society Edinburgh*, v. 45, p. 223–233.
- Reid, C.D., Maherali, H., Johnson, H.B., Smith, S.D., Wullschlegel, S.D., Jackson, R.B., 2003, On the relationship between stomatal characters and atmospheric CO₂: *Geophysical Research Letters*, v. 30, no. 19, doi: 10.1029/2003GL017775.
- Royer, D.L., 2001, Stomatal density and stomatal index as indicators of paleoatmospheric CO₂ concentration: Review of Palaeobotany and Palynology, v. 114, p. 1–28, doi: 10.1016/S0034-6667(00)00074-9.
- Smith, W.K., and Donahue, R., 1991, Simulated effect of altitude on photosynthetic CO₂ uptake potential in plants: *Plant, Cell & Environment*, v. 14, p. 133–136.
- Terashima, I., Masuzawa, T., Ohba, H., and Yokoi, Y., 1995, Is photosynthesis suppressed at higher elevations due to low CO₂ pressure: *Ecology*, v. 76, p. 2663–2668.
- von Caemmerer, S., and Farquhar, G.D., 1981, Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves: *Planta*, v. 153, p. 376–387, doi: 10.1007/BF00384257.
- Woodward, F.I., and Bazzaz, F.A., 1988, The responses of stomatal density to CO₂ partial-pressure: *Journal of Experimental Botany*, v. 39, p. 1771–1781.