

Overall Story

- Accelerating charges generate free space **electromagnetic waves**.
- Light is an EM wave and can be **reflected** and **refracted**.
- **Mirrors** and **lenses** form images of objects by collecting rays emitted from an object through reflection or refraction.

Maxwell's Equations

$$\oint \vec{E} \cdot d\vec{A} = \frac{Q}{\epsilon_0}$$

Gauss' Law

$$\oint \vec{B} \cdot d\vec{A} = 0$$

Gauss' Law for Magnetism – no magnetic monopoles

$$\oint \vec{E} \cdot d\vec{s} = - \frac{d\Phi_B}{dt}$$

Faraday's Law


$$\oint \vec{B} \cdot d\vec{s} = \mu_0 I + \mu_0 \epsilon_0 \frac{d\Phi_E}{dt}$$

Ampère-Maxwell Law

$$\vec{F} = q\vec{E} + q\vec{v} \times \vec{B}$$

Lorentz Force Law

EM Waves



$$E = E_{\max} \cos(kx - \omega t)$$

$$B = B_{\max} \cos(kx - \omega t)$$

$$\frac{E}{B} = \frac{E_{\max}}{B_{\max}} = \frac{\omega}{k} = c$$

$$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$$

Speed of Light

$$\omega = 2\pi f$$

Angular Frequency

$$\lambda = \frac{c}{f}$$

Wavelength

$$k = \frac{2\pi}{\lambda}$$

Wavenumber


$$\frac{\omega}{k} = c$$

Energy Carried by EM Waves

$$\vec{S} = \frac{1}{\mu_0} \vec{E} \times \vec{B}$$

Poynting Vector (W/m²)

$$S = \frac{EB}{\mu_0} = \frac{E^2}{\mu_0 c} = \frac{c}{\mu_0} B^2$$



$$I = S_{av} = \frac{E_{max} B_{max}}{2\mu_0} = \frac{E_{max}^2}{2\mu_0 c} = \frac{c}{2\mu_0} B_{max}^2$$

$$u_E = u_B = \frac{1}{2} \epsilon_0 E^2 = \frac{B^2}{2\mu_0}$$


$$u = u_E + u_B = \epsilon_0 E^2 = \frac{B^2}{\mu_0}$$

$$u_{av} = \epsilon_0 (E^2)_{av} = \frac{1}{2} \epsilon_0 E_{max}^2 = \frac{B_{max}^2}{2\mu_0}$$


$$I = S_{av} = cu_{av}$$

Momentum and Radiation Pressure

	Momentum	Pressure
Complete Absorption	$p = \frac{U}{c}$	$P = \frac{F}{A} = \frac{1}{A} \frac{dp}{dt} = \frac{S}{c}$
Complete Reflection	$p = \frac{2U}{c}$	$P = \frac{2S}{c}$



Reflection and Refraction





$$\theta'_1 = \theta_1$$

$$n = \frac{c}{v}$$

$$n_1 \sin \theta_1 = n_2 \sin \theta_2$$


$$\sin \theta_c = \frac{n_2}{n_1}$$

Dispersion and Prisms



$$n = n(\lambda)$$

Images by Mirrors



$$p = q$$

$$M = 1$$

Front, or
real, side

Back, or
virtual, side

p and q positive

Incident light


p and q negative

Reflected light

p and q negative

No light


Convex or concave mirror




$$\frac{1}{p} + \frac{1}{q} = \frac{1}{f}$$

$$f = \frac{R}{2}$$


$$M = \frac{h'}{h} = -\frac{q}{p}$$



Images by Refraction and Lenses




$$\frac{n_1}{p} + \frac{n_2}{q} = \frac{(n_2 - n_1)}{R}$$



$$\frac{1}{p} + \frac{1}{q} = \frac{1}{f}$$

$$\frac{1}{f} = (n-1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$


$$M = \frac{h'}{h} = -\frac{q}{p}$$



Front

p positive
 q negative

Incident light








Back

p negative
 q positive


Refracted light

Ray Diagrams



Optical Instruments


The Camera



$$I \propto D^2/f^2$$


$$f\text{-number} = \frac{f}{\#} = \frac{f}{D}$$

Magnifier



$$m_\theta \approx \frac{25\text{cm}}{f}$$

Compound Microscope



$$M = -\frac{p_1}{f_o} \frac{25\text{cm}}{f_e}$$